

1. **Diphyrama singularis.** (Tab. V. fig. 12.)*Diphyrama singularis*, Bates, Trans. Ent. Soc. 1872, p. 188.*Hab.* NICARAGUA, Chontales (*Belt*).

LISTROPTERA.

Listroptera, Serville, Ann. Soc. Ent. Fr. 1834, p. 71.

Eight species have been described of this genus, all from Tropical South America. One of them extends its range far into Central America.

1. **Listoptera aterrima.***Listoptera aterrima*, Germar, Ins. Spec. Nov. p. 497 ; Bates, Trans. Ent. Soc. 1872, p. 188².

Hab. NICARAGUA, Chontales (*Belt*², *Janson*) ; COSTA RICA (*Rogers*).—SOUTH AMERICA to Santa Catharina, SOUTH BRAZIL.

DIHAMMOPHORA.

Dihammophora, Chevrolat, Arcana Naturæ, p. 51 (1859) ; Lacord. Gen. Col. ix. p. 108.*Listroptera* (partim), White, Cat. Long. Col. B. M. ii. p. 207.

Also an exclusively American genus, ranging in the south as far as Buenos Ayres, and in the north to the *tierras calientes* of Mexico. Sixteen species have been described.

1. **Dihammophora chontalensis.** (Tab. V. fig. 13.)*Dihammophora Chontalensis*, Bates, Trans. Ent. Soc. 1872, p. 188.*Hab.* NICARAGUA, Chontales (*Belt*).2. **Dihammophora aurovittata.**

D. marginicollis similis, at thoracis vittis sericeo-pubescentibus ; linearis, postice vix dilatata, atra ; frontis marginibus auro-sericeis ; thorace lineari, dorso atro, vittis duabus (marginem posticum vix attingentibus) auro-pubescentibus sericeis (vitta dorsali nigra nuda grosse punctata), subtus fulvo ; elytris crebre sublineatim punctatis, lateribus anguste bicostatis ; pedibus nigris, nitidis, femoribus basi flavo-testaceis.

Long. 3 lin.

Hab. GUATEMALA, Capetillo, Zapote (*Champion*).3. **Dihammophora dispar.***Dihammophora dispar*, Chevr. Arc. Nat. p. 52.*Hab.* MEXICO.

RHOPALOPHORA.

Rhopalophora, Serv. Ann. Soc. Ent. Fr. 1834, p. 100 ; Chevr. Arc. Nat. p. 57.*Tinopus*, Leconte, Journ. Ac. Phil. ser. 2, ii. p. 19.

Twenty-five species of this elegant genus have been described, all American. Although most numerous in the tropical region, the genus extends into temperate