

8. **Rhopalophora versicolor.** (Tab. V. fig. 15.)

Rhopalophora versicolor, Chevr. Arc. Nat. p. 62¹; Bates, Trans. Ent. Soc. 1872, p. 188².

Hab. NICARAGUA, Chontales (*Belt*², *Janson.*)—SOUTH AMERICA, Colombia¹.

9. **Rhopalophora serripes.**

Rhopalophora serripes, Bates, Trans. Ent. Soc. 1872, p. 188.

Hab. NICARAGUA, Chontales (*Belt*).

COSMISOMA.

Cosmisoma, Serville, Ann. Soc. Ent. Fr. 1834, p. 19; Lacord, Gen. Col. ix. p. 112.

Cosmosoma, Har. & Gemm. Cat. Col. p. 2948.

One of the most beautiful genera of American Longicornia, belonging to the Rhopalophorini group, and distinguished by the brushes of silky hairs on the antennæ, which are present, of great diversity of shape and colour, on the fifth joint, but sometimes in addition on other joints, and even on the hind tibiæ. Nineteen species are known, from South Brazil to Nicaragua.

1. **Cosmisoma martyra.** (Tab. V. fig. 16.)

Cosmisoma martyra, Thoms. Class. Long. p. 180¹.

Cosmisoma martyra, Bates, Trans. Ent. Soc. 1872, p. 189².

Hab. GUATEMALA, Zapote (*Champion*); NICARAGUA, Chontales (*Belt*²); COSTA RICA¹.

The single specimen from Guatemala (a female) is of a brighter brassy-green general colour than the Nicaraguan examples, and the thorax is a little less closely punctulated.

2. **Cosmisoma titania.** (Tab. V. fig. 17.)

Cosmisoma Titania, Bates, Trans. Ent. Soc. 1870, p. 407¹.

Hab. NICARAGUA, Chontales (*Belt*¹, *Janson*).

3. **Cosmisoma plumicornis.**

Leptura plumicornis, Drury, Ill. iii. p. 74, t. 49. f. 3.

Cosmisoma plumicornis, Pascoe, Trans. Ent. Soc. 3rd ser. v. p. 291¹.

Hab. COSTA RICA?—SOUTH AMERICA, Colombia¹.

I obtained a specimen of this beautifully ornamented species out of a small collection of insects received by Mr. Gerrard from an ornithological traveller, and obtained, he believed, from Costa Rica.

OZODES.

Ozodes, Serville, Ann. Soc. Ent. Fr. 1834, p. 98.

Ten species have been recorded of this distinct genus, peculiar to Tropical America.