

1. **Ozodes multituberculatus.**

Ozodes multituberculatus, Bates, Tr. Ent. Soc. 1870, p. 409¹.

Hab. NICARAGUA, Chontales (*Belt*¹, *Janson*).

2. **Ozodes xanthophasma.** (Tab. VI. fig. 14.)

Ozodes xanthophasma, Bates, Tr. Ent. Soc. 1872, p. 189¹.

Hab. NICARAGUA, Chontales (*Belt*¹).

ORNITHIA.

Ornithia, Thomson, Syst. Céramb. p. 213; Lacord. Gen. Col. ix. p. 117.

Trichophorus, Guér. Icon. R. A. texte, iii. p. 228.

Ozodes, Sturm.

The following is the only known species:—

1. **Ornithia chevrolati.**

Trichophorus Chevrolatii, Guér. Icon. R. A. texte, iii. p. 228¹.

Ozodes mexicanus, Sturm, Cat. ed. 1843, t. 6. f. 4, 5.

Ornithia Chevrolatii, Bates, Trans. Ent. Soc. 1872, p. 190³.

Hab. MEXICO¹, Misantla (*Höge*); HONDURAS (*Blancaneaux*); NICARAGUA, Chontales (*Belt*³, *Janson*).

CHRY SOPR ASIS.

Chrysoprasis, Serville, Ann. Soc. Ent. Fr. 1834, p. 5.

Callichroma, Germar, Ins. Spec. Nov. p. 496.

A genus of small metallic-coloured Cerambycidae, differing from *Callichroma*, to which they have a superficial resemblance in colours, by the closed sockets of the middle coxæ and other characters. The species are very numerous in the equatorial forest-region of America, and some are found as far south as the La Plata; but they diminish in numbers as we advance northward, and none have yet been recorded from Mexico. About fifty species are known, all American.

1. **Chrysoprasis belti.** (Tab. VI. fig. 2.)

Chrysoprasis Belti, Bates, Trans. Ent. Soc. 1872, p. 190.

Hab. NICARAGUA, Chontales (*Belt*).

2. **Chrysoprasis sthenias.**

Chrysoprasis sthenias, Bates, Trans. Ent. Soc. 1870, p. 411.

Var. Femoribus nigris, vix æneo tinctis.

Hab. NICARAGUA, Chontales (*Belt*).—SOUTH AMERICA, Amazons.

A robust species with short spined antennæ, reticulate-punctate pronotum and metasternum, and thickly punctured femora. South-American specimens have brassy-green