

Colobothea chontalensis (p. 188).

To the locality given, add:—PANAMA, Volcan de Chiriqui, Bugaba (*Champion*).

7 (A). **Colobothea aleata**. (Tab. XXV. fig. 10.)

C. hebraica proxime affinis, sed minor et gracilior; elytris guttis multo numerosioribus albis subtessellatis haud in fasciis tribus congestis; thorace medio lineolis tribus, 1° anteriori, 2° et 3° posterioribus, his quam in *C. hebraica* magis approximatis.

Segmento abdominali apicali, ♂ ♀, simplici, sicut in *C. hebraica*.

Long. $4\frac{1}{2}$ –6 lin.

Hab. NICARAGUA, Chontales (*Belt*); PANAMA, Volcan de Chiriqui, Bugaba (*Champion*).

Colobothea distincta (p. 189).

To the localities given, add:—GUATEMALA, Mirandilla (*Champion*); PANAMA, Bugaba (*Champion*).

13. **Colobothea** — ?

Hab. PANAMA (*Boucard*).

A single example in very dilapidated condition of what appears to be a very distinct species.

14. **Colobothea** — ?

Hab. PANAMA, Chiriqui (*Trötsch*).

A single worn specimen, distinct from any of the Central-American species recorded here, and apparently allied to *C. latevittata*, Bates.

CARNEADES (p. 190).

Carneades hemileuca (p. 190).

To the locality given, add:—PANAMA, Bugaba (*Champion*).

6. **Carneades championi**. (Tab. XXV. fig. 11.)

C. superbæ et *C. grandis* proxime affinis et similis, differt thorace nebula discoidali transversa nigro-fusca (lateribus haud maculatis) elytrisque macula communi suturali paullo post scutellum, alteris utrinque ante medium quatuor, fascia recta post medium maculaque transversa quadrata ante apicem, nigris. Cretaceo-alba, interdum ochraceo-fulva; antennis nigris, articulis basi griseo-albis; pedibus ventrequae sicut in *C. superba* variegatis.

Long. 6–13 lin. ♂ ♀.

Hab. PANAMA, Volcan de Chiriqui, Bugaba (*Champion*).

Differs from *C. superba* and *C. grandis* only in the number and position of the black markings; but in these differences the large series of individuals exhibit unwavering constancy. Besides the absence of distinct black spots from the thorax, the chief