

A single example from Chiriqui, with which a specimen received as from "Brazil" agrees in every respect, except that the ashy-grey adpressed pubescence is not so clearly defined and the general colour more castaneous, owing doubtless to immaturity.

2. *Bactriola paupercula*.

Fusca (elytris subaenescens), cinereo-fusco tenuiter pubescens, thorace utrinque roseo; antennis nigris, articulis 3^o-5^m basi cinereis; pedibus rufo-testaceis, femoribus basi, tibiis apice tarsisque, nigro-fuscis.

Long. 2½ lin.

Hab. PANAMA, Volcan de Chiriqui (*Champion*).

Several examples.

DRYCOTHEA (p. 193).

Drycothea sallæi (p. 193).

To the localities given, add:—BRITISH HONDURAS (*Blancaneaux*); GUATEMALA, Cahabon, Las Mercedes, El Reposo, El Tumbador (*Champion*).

This species differs from *D. stictica*, which it very much resembles in colour, by its shorter form and relatively larger, especially broader, thorax. More definite points of difference are the small lower lobe of the eyes, narrowed below, and the slightly convex mesosternum in *D. sallæi*; the lower lobe of the eyes being very large and broad, as broad or broader below than above, and rounded on the inner side, and the mesosternum gibbous, in *D. stictica*.

Drycothea stictica (p. 193).

To the localities given, add:—MEXICO, Jalapa (*Höge*); BRITISH HONDURAS (*Blancaneaux*); PANAMA, Volcan de Chiriqui, Bugaba (*Champion*).

Var. *D. lepidiota*. Oculi lobo inferiori paullo minus rotundato; mesosterno mediocriter convexo.

Hab. MEXICO, Toxpam (*Sallé*).

Labelled *Estola lepidiota*, Chevr. (MS.), in the Sallé collection.

Same habits as *Estola* (*Champion*).

Drycothea cribrata (p. 194).

To the locality given, add:—MEXICO, Jalapa (*Höge*).

5. *Drycothea spreta*.

D. sticticæ valde affinis et similis; minor et angustior, densius fulvo vel fusco-cinereo incumbenti-pubescens et magis confertim et subtilius (apud elytra confusus) punctata. Elongato-oblongo-linearis; oculis lobo inferiori magno, oblongo, sed minus rotundato; thoracis spina parva, acuta; elytris haud perspicue striato-punctatis, confuse, hic illic subseriatim, punctulatis, interstitiis haud convexis; mesosterno supra plano, antice rotundato-declivi.

Long. 4 lin.

Hab. MEXICO, Cordova (*Sallé*, *Höge*), Jalapa (*Höge*).

This species seems to be the common form in Mexico, and to be connected