

Hab. PANAMA, Volcan de Chiriqui (*Champion*).

One example only.

14. *Amphionycha pulchricollis*. (Tab. XXV. fig. 19.)

Sublinearis, nigra; elytris (apice epipleurisq. nigro-fuscis exceptis) clare fulvo-ochraceis, occipite vittis duabus antice convergentibus vittaque utrinque thoracis laterali et linea dorsali pallidius ochraceis; antennis (articulis 1° et 2° nitide nigris exceptis) pedibusque (tarsis apiceque tibiis nigris exceptis) melleo-flavis; capite robusto, inter antennis mediocriter concavo; thorace cylindrico, basi sat constricto; elytris apice sinuato-truncatis, utrinque bidentatis, dente exteriori multo longiori, epipleuris altis striato-punctatis bicarinatis (carina inferiori antice obsoleta), dorso punctulato-striato. Antennis tenuibus sicut in *A. pluricostata* et *A. bicarinata* (♂) brevioribus, subtus ciliatis, articulis versus apicem multo abbreviatis.

Long. $4\frac{3}{4}$ lin. ♂.

Hab. GUATEMALA, San Gerónimo (*Champion*).

One example only.

15. *Amphionycha ocreata*.

Sat anguste cylindrica, nigro-fusca, erecte pilosa; fronte thoraceque subtiliter albo-cinereo-tomentosis, hoc dorso posteriori plaga magna nigro-fusca; elytris sutura fulvo-ochracea, metathoracis episternis ventrequa rufo-testaceis, pedibus rufo-testaceis, femoribus apice tibiisque (posticis apice exceptis) nigris nitidis; antennis tenuibus, subtus longe parum dense ciliatis (articulo 4° valde elongato), nigris, scapo subtus rufo-testaceo, articulis 3° apice, 6° basi, 4° et 5° totis, albo-testaceis; capite inter antennis concavo; thorace cylindrico; elytris apice obtuse truncatis, epipleuris altis striato-punctatis carina superiori valida inferiori solum versus apicem elevata, dorso striato-punctulato.

Long. $4\frac{1}{4}$ lin.

Hab. PANAMA, Volcan de Chiriqui (*Champion*).

Similar in colours and shape to *Isomerida lineata*, ♂, from which its generic characters—the slenderer antennæ with relatively much longer third and fourth joints—amply distinguish it. It also resembles *A. pubicornis*, but is longer and narrower than that species.

ALAMPYRIS (p. 218).

Alampyris marginella (p. 220).

To the Mexican locality given, add:—Cerro de Plumas (*Höge*).

9. *Alampyris cretaria*.

Elongato-oblongo-ovata, breviter erecte pilosa, fusco-nigra; thorace utrinque plaga magna quadrata, densissime et subtilissime cretaceo-albo tomentoso; capite albo-tomentoso, vittis duabus frontalibus altera utrinque post oculus nigro-fuscis; elytris sutura et margine fulvis; antennis brevibus, scapo elongato gradatim clavato, articulo 3° quam scapus haud longiori, 5°–11^m valde abbreviatis, rufo-testaceis, articulis apice, scapo et articulis 8°–11^m toto, nigro-fuscis, scapo undique setoso, 2°–4^m subtus longe ciliatis; thorace antice angustato, post medium lateribus paullo tumidis; elytris mediocriter convexis, epipleuris nec verticalibus nec carinatis, apice singulatim subacuminatim rotundatis, dorso confuse punctulatis; ventre (♂?) immaculato; femoribus anticis rufo-testaceis.

Long. 6 lin.

Hab. MEXICO (ex coll. A. Turner). Two examples, apparently males.