

colore supra sicut in *L. unicolori* toto isabellino; antennis nigris (scapo piceo), articulis 5<sup>o</sup>-9<sup>m</sup> pallide testaceis; elytris mox post humeros (imprimis valde gradatim) explanato-dilatatis, prope suturam solum punctulatis utrinque bicostatis; abdomine pedibusque pubescentibus.

Long.  $6\frac{1}{2}$  lin.

*Hab.* PANAMA, Bugaba (*Champion*).

This species and *L. mimica* form a subgroup, in which the antennal scape is considerably longer than in the rest of the genus. The dilatation of the elytra is rather wider and more gradual from the shoulder to near the apex than it is in *L. mimica* and *Lycaneptia amicta*.

#### 4. *Lycidola eximia*.

Supra aurantiaca, elytris fascia recta versus basin alteraque apicali, nigris; antennis nigris, articulis 3<sup>o</sup> et 4<sup>o</sup> sicut in *L. unicolori* æqualibus; vertice thoraceque vitta mediana indistincta fusca; elytris paullo post humeros sat late explanato-dilatatis haud angulatis, undique confertim punctatis utrinque quadricostulatis; pectoris lateribus, ventre, tibiis et tarsis fusco-nigris, pectore medio femoribusque fulvo-testaceis.

Long. 5 lin.

*Hab.* PANAMA, Bugaba (*Champion*).

One example only.

### ADDENDA.

TETROPIUM (to follow the genus *Spondylis*, pp. 14, 239).

*Tetropium*, Kirby, Fauna Bor.-Amer. iv. p. 174; Lacordaire, Gen. de Col. viii. p. 208.

A genus hitherto known only from the higher temperate latitudes of America, Europe, and Asia, whence five species have been described.

#### 1. *Tetropium opacipenne*.

*T. aulico* relative multo longius, nigrum, fusco-pilosum; elytris opacis adpresso-pubescentibus, femoribus rufis; thorace fere sicut in *T. aulico* rotundato, confertim punctulato, dorso callis duobus lineolaque dorsali paullo minus dense punctatis; elytris costis vix perspicuis.

Long.  $7\frac{1}{2}$  lin. ♀.

*Hab.* MEXICO, Jacale, Orizaba (*Sallé*).

One example.

#### 2. *Tetropium auripilis*.

Præcedenti quoad formam simile. Nigrum, nitidum, pilosum; elytris cinnamomeis sericeo-aureo-pubescentibus; thorace (♂) paullo longiori, medio subangulatim dilatato, confertim punctato, dorso tricalloso; scutello nigro; elytris costis inconspicuis; pedibus nigris, nitidis, tarsis piceis.

Long.  $7\frac{1}{2}$  lin. ♂.

*Hab.* MEXICO, Las Vigas (*Flohr, coll. Bates*).

One example.