

Hab. GUATEMALA, near the city, Capetillo (*Champion*).

Antennæ only moderately long and broad; third and fourth joints slender, subequal, the penultimate joint about as long as broad. Head small; eyes scarcely at all prominent, widely separated. Thorax rather short, rounded at the sides so as to be much narrowed in front, black, very finely punctate, but with some larger, but still quite fine, punctures scattered over it, slightly shining, the surface receiving a somewhat leaden tint from the very minute pubescence; basal lobe foveolate in front of the scutellum. Elytra rather finely striate, each with a large red spot occupying the larger part of the surface, reaching the outer margin, but not the base, apex, or suture; interstices densely, minutely punctulate, the pubescence very minute. Pygidium broad, rather large, the pubescence very fine. Posterior femora not dilated, each with a rather slender acuminate tooth; tibiæ straight, only moderately broad, without mucro.

Only one example was met with in each locality; they represent the two sexes, the last ventral ring in the male being considerably excised. Although superficially similar to some of the maculate varieties of *B. instabilis*, this species is very distinct therefrom; the front coxæ are separated by a fine lamina, the hind femora not canaliculate.

64. *Bruchus trisignatus*.

Brevis, subrotundatus, minus convexus, niger, omnium brevissime pallido vestitus, elytro singulo macula magna humerali rufa; antennarum basi testacea; prothorace densissime punctulato; pygidio basi albido trimaculato; femoribus posterioribus dente singulo armatis.

Long. 3 millim.

Hab. PANAMA, Bugaba (*Champion*).

Head short and broad; eyes widely separated. Antennæ short and broad, the penultimate joints transverse. Thorax broad, rather short, greatly narrowed in front, the sides rounded; surface dull owing to a very dense minute punctuation, the basal lobe not impressed. Elytra each with a large humeral red spot occupying nearly half the surface, rather finely striate, the striæ punctured, the interstices densely punctulate. Pygidium broad, punctate, with three very conspicuous large spots of white pubescence at the base. Hind femora scarcely incrassate, each armed with a rather small tooth; tibia short, very thick, straight, with a short, strong mucro at the extremity. Front coxæ distinctly separated.

Only one example was obtained of this very distinct species.

65. *Bruchus cruentatus*.

Bruchus cruentatus, Horn, Trans. Am. Ent. Soc. iv. (1873), p. 325¹.

Hab. NORTH AMERICA, Georgia, Illinois, Dacota, and Texas¹.—MEXICO, Guanajuato (*Sallé*), Jalapa (*Höge*).