

66. *Bruchus flaviventris*.

Brevis, niger, supra ochraceo, subtus pallido-griseo, vestitus; antennis, pedibus abdomineque flavis; elytris disco rufo bimaculato; femoribus posterioribus muticis.

Long. $2\frac{1}{3}$ millim.

Hab. GUATEMALA, Dueñas (*Champion*).

Antennæ pale yellow, short, and very broad; third joint rather smaller than the second; joints six to ten strongly transverse. Eyes rather small, widely separated. Thorax transverse-conic, the sides scarcely rounded in front; surface very dull, densely sculptured, with larger punctures in addition to the dense sculpture, the pubescence very pallid ochraceous, uniform and rather abundant. Elytra moderately deeply striate, clothed with pubescence like the thorax, each with a large pale red spot near the suture. Pygidium yellow. Legs including the front coxæ and the hind tarsi yellow. Hind femora not dilated, but rather short and broad, unarmed; tibiæ quite destitute of mucro.

Of this very distinct species, only one example has been procured; though it bears a superficial resemblance to *B. cruentatus*, it is not closely allied to it or to any other recorded here.

67. *Bruchus medialis*.

Brevis, convexus, niger, supra ochraceo, subtus pallido-griseo, vestitus; antennis pedibusque testaceis, illis articulis sexto ad decimum fuscis; elytrorum medio late rufo-fasciato; femoribus posterioribus dente parvo armatis.

Long. 2 millim.

Hab. MEXICO, Guanajuato (*Sallé*).

Antennæ short and broad, third and fourth joints small, each smaller than the second joint. Eyes rather small and widely separated. Thorax short, greatly narrowed in front, the sides rounded in front, the surface densely and rather finely punctulate, and with a rather dense yellow pubescence. Elytra short, rather deeply striate, the interstices aciculate not punctulate; each with a very large pale-red mark nearer the base than the apex, starting from the outer margin and reaching the suture, except for the sutural interval which remains black. Pygidium rather abundantly clothed with pallid pubescence, rather large and convex. Legs bright red, front coxa and the base of the hind femora black. Hind femora not dilated, and armed only with a small sharp tooth; tibia with the anterior angle scarcely longer than the other digits.

Of the three examples on which I describe this species two are males, and have the terminal ventral ring shorter in the middle.

68. *Bruchus interruptus*.

Brevis, minus convexus, subrotundatus, niger, breviter albido pubescens; antennis pedibusque testaceis, illis extrorsum fuscescentibus; elytris rufo-signatis, striatis, striis fortiter punctatis; femoribus posterioribus dente sat elongato armatis.

Long. $2\frac{1}{3}$ millim.