

21. *Lema trilineata*.

Lema trilineata, Oliv. Ent. vi. 1808, p. 739, t. 2. f. 20; Lacord. Mon. p. 418; Harris, Rep. Ins. Massach. injur. p. 95.

Lema, var. *immaculicollis*, Chevr. Col. Mex. cent. ii. 1835, no. 112.

Lema trivirgata, Leconte, Col. of Kansas, 1859, p. 22.

Lema lecontei, Clark, Cat. Phyt. 1866, p. 31.

Lema trivittata, Say, Journ. Ac. Phil. iii. p. 429.

Hab. N. AMERICA.—MEXICO; HONDURAS; COSTA RICA.—CUBA.

Var. *immaculicollis*, Chevr., has also been received from Honduras.

22. *Lema nigrovittata*.

Lema nigrovittata, Guér. Ic. Règn. Anim. iii. p. 262¹; Lacord. Mon. p. 421.

Lema melanocephala, Dej. Cat. 3rd ed. p. 386.

Hab. N. AMERICA, California¹.—MEXICO.

This species is very closely allied to the preceding one, and is equally variable. It is, however, less robust and smaller; and the submarginal stripe is not placed so close to the outer margin as in *L. trilineata*.

23. *Lema opulenta*.

Lema ornata, Baly, Ann. & Mag. Nat. Hist. ser. 3, xvi. p. 158¹ (nec Gravenh.).

Lema opulenta, Gemm. & v. Har. Cat. Col. xi. p. 3258².

Hab. GUATEMALA¹, Zapote, Dueñas (*Champion*).

The Munich Catalogue name, *L. opulenta*, Harold, has been substituted for that given by Mr. Baly, whose title is unfortunately preoccupied. I am, however, quite unable to find any reference to the subject in part xiii. of the Coleopterologische Hefte, or any mention of the species.

24. *Lema abrupta*.

Lema abrupta, Lacord. Mon. p. 422.

Hab. MEXICO.

25. *Lema salvini*.

Lema salvini, Baly, Cistula Entom. pt. ii. 1878, p. 313.

Hab. GUATEMALA, near the city (*Salvin*).

26. *Lema semisepta*.

Lema semisepta, Lacord. Mon. p. 423¹.

Hab. GUATEMALA, Capetillo (*Champion*).—BRAZIL¹.

I refer a single specimen sent from Guatemala to this species; it differs, however, in several particulars: the tibiæ and knees are black, as well as the antennæ, with the exception of the last two joints, which are rufous; the thorax is as broad as long; and the elytral spot is absent as in Lacordaire's var. *a*.