

**16. *Mastostethus versicolor*.**

*Mastostethus versicolor*, Lacord. Mon. p. 662.

*Hab.* MEXICO.

**17. *Mastostethus phaleratus*.**

*Mastostethus phaleratus*, Klug, Jahrb. Ins. p. 221 ; Lacord. Mon. p. 663.

*Mastostethus dohrnii*, Baly, Journ. Ent. i. p. 281, t. 13. f. 5.

*Hab.* MEXICO, Oaxaca ; COSTA RICA, Caché (*Rogers*).

As will be seen by the above synonyms, I have united Mr. Baly's species with that of Klug ; from Lacordaire's description I am absolutely unable to distinguish them. Curious mistakes, however, seem to have taken place in regard to this insect, whose elytra are described by Mr. Baly as "clothed with very short suberect hairs;" in the type before me I am unable, even with the strongest glass, to discover any hairs. In the catalogue of Gemminger and von Harold, Mr. Baly's species is classified in the genus *Agathomerus*, no doubt on account of the description in regard to the pubescence of the elytra ; while Mr. Baly himself has described it as belonging to *Mastostethus*. In every respect the insect agrees with the description in Lacordaire's Monograph.

About twelve specimens have been collected by Mr. Rogers ; and they all agree perfectly with the type, and show no trace of hairs on the elytra. It is a true *Mastostethus*.

**18. *Mastostethus hieroglyphicus*.**

*Mastostethus hieroglyphicus*, Klug, Jahrb. Ins. p. 222 ; Lacord. Mon. p. 663.

*Hab.* MEXICO.

**19. *Mastostethus novemmaculatus*. (Tab. I. fig. 19.)**

*Mastostethus novemmaculatus*, Klug, Jahrb. Ins. p. 211 ; Chevr. Col. Mex. Cent. ii. ; Lacord. Mon. p. 668.

*Hab.* MEXICO.

The antennæ of this species (the colour of which Lacordaire could not give, on account of his specimen only possessing one joint) are entirely rufous, and slightly longer than the thorax.

**20. *Mastostethus placidus*. (Tab. I. fig. 22.)**

*Mastostethus placidus*, Baly, Cist. Ent. i. p. 127.

*Hab.* MEXICO.

**21. *Mastostethus lavatus*.**

*Mastostethus lavatus*, Baly, Journ. Ent. i. p. 281, t. 13. f. 6.

*Hab.* MEXICO, Oaxaca.