

specimen, and seems scarcely to vary from most of the others belonging to his fourth section.

### 11. *Pachybrachys incrassatus*.

*Pachybrachys incrassatus*, Suffr. Monogr. vii. p. 206<sup>1</sup>.

*Hab.* MEXICO, Jalapa<sup>1</sup>; GUATEMALA, San Gerónimo (*Champion*).

The shorter antennæ (which do not extend to half the length of the body), the close and strong punctuation of the thorax, and the irregularly sinuate costæ of the elytra form the principal distinguishing characters of this species, to which I refer a male (a sex unknown to Suffrian) sent by Mr. Champion, and which agrees well enough with the type in the Berlin Museum. The general colour is yellow, with brown lines and punctures.

### 12. *Pachybrachys longulus*.

*Pachybrachys longulus*, Suffr. Monogr. vii. p. 207.

*Hab.* MEXICO.

Again a closely allied species to the preceding ones, but separated by the long and narrow cylindrical body and the white colour at the sides of the abdominal segments. The type in the Berlin Museum is a female, the antennæ of which reach nearly to the end of the body. The elytra are very closely and irregularly covered with brown punctures; and the legs and underside are of the same colour.

### 13. *Pachybrachys punctatissimus*. (Tab. IV. fig. 20.)

Broad, parallel, brown below; head and thorax closely punctured, the latter with two longitudinal bands and the margins yellow; elytra light yellow, covered closely with brown punctures.

Length  $1\frac{3}{4}$ –2 lines.

Head exceedingly closely rugose punctate, brown, with irregular spots of yellow; antennæ extending to one third the length of the body, the apical joints rather robust and much widened, black, basal joint fulvous; thorax of equal width, its sides straight, posterior margin distinctly produced at the middle; surface very closely, evenly, and deeply punctate, brown, interrupted here and there by small yellow spots, and divided longitudinally by two bands of the same colour, more or less distinct; scutellum dark brown; elytra punctured like the thorax, the punctures themselves brown, the interstices light yellow, and now and then arranged into smooth, narrow, longitudinal costæ; underside and legs darker or lighter brown; female with some brown spots near the apex of the elytra.

*Hab.* MEXICO, La Parada, Cuernavaca (*Sallé*).

A species allied to *P. longulus*, Suffr., from which it is distinguished by its shorter body, antennæ, and the very close and rugose punctuation of the head; that of the elytra in *P. longulus* is also finer, and does not extend quite to the apex; and the abdomen in the present species is without the white spots. A specimen in the collection of Mr. Baly slightly exceeds two lines, and is proportionally robust.