

piceous; antennæ half the length of the body, black, the basal joint fulvous, the third and following joints rather slender and each longer than broad; thorax rather longer than broad, the sides somewhat deeply constricted, the basal sulcation obsolete, the surface obscure fulvous, with some nearly obsolete longitudinal black spots at the sides, the disc impunctate; scutellum fulvous; elytra convex, subcylindrical, without any trace of a basal depression, the punctuation only well-marked anteriorly near the suture, the rest of the disc very finely and obsoletely punctured, the interstices flat; underside fulvous, the knees, the apical third of the tibiæ, the tarsi, and a small spot at the side of each abdominal segment, black.

Hab. PANAMA, David in Chiriqui (*Champion*).

A single specimen. On account of the filiform (not widened) joints of the antennæ I am unable to treat this insect as a variety of *L. varipes* or *L. nupta*; the upper surface immaculate, and the apices only of the tibiæ (instead of the middle portion, as in *L. intricata*) are black. In the sculpture and shape of the elytra this species very closely resembles *L. intricata*.

36 (E). *Lema modesta*.

Pale fulvous; labrum piceous; antennæ with shortened terminal joints; thorax longer than broad, impunctate; elytra convex, distinctly punctured towards the base only, the rest of the surface very finely punctate; underside and legs obscure fulvous, the abdomen lighter.

Length $2\frac{1}{2}$ lines.

Hab. PANAMA, Bugaba (*Champion*).

This insect is closely allied to *L. immaculipennis*, but I believe distinct; it differs thus:—the antennæ are entirely fulvous, and the terminal joints (instead of being elongate) are short and gradually, but slightly, thickened; the thorax is longer, and the underside and legs are without black spots.

The elytra are shaped and sculptured exactly as in *L. immaculipennis*.

A single specimen only was obtained.

Lema atricornis (p. 9).

To the locality Mexico, add:—Playa Vicente (*Sallé*).

Lema sexnotata (p. 9).

To the locality Mexico, add:—Jalapa, Cordova (*Höge*), Vera Cruz (*Sallé*).

39 (A). *Lema intricata*.

Fulvous, the antennæ, four spots on the head, a spot on each side of the thorax, and the scutellum, black; elytra strongly punctate-striate, a spot at the sides before and another below the middle, and a sutural spot below the base, black; breast with two spots; legs black at the base, at the knees, and at the middle of the tibiæ.

Length $2\frac{1}{2}$ lines.

Head impunctate, fulvous, the space round the eyes punctured, the vertex with a central black spot, the base with three spots placed transversely; antennæ less than half the length of the body, black, the basal joint fulvous; thorax not longer than broad, deeply constricted at the sides, the basal groove rather deep, the surface with a few minute punctures at the middle of the disc, the sides with a black spot at the middle