

varies considerably in size; and the scutellum is clothed with white hairs (not smooth as in *E. pilatei*). Lacordaire describes the second to the fifth joints of the antennæ as fulvous; in all the specimens I have examined the second and third joints only are of that colour.

5. *Euryscopa nigratarsis*. (Tab. XXXVII. fig. 1.)

Broadly subcylindrical, black; legs fulvous, the tarsi black; thorax sparingly punctured on the disc; elytra distinctly punctured anteriorly, very finely so below the middle, the anterior third and a round spot near the apex fulvous.

Length $2\frac{1}{2}$ lines.

Hab. PANAMA, Bugaba (*Champion*).

E. nigratarsis, of which a single female specimen only is before me, seems closely allied to *E. carnifex* and *E. specularis*, Lac., without agreeing sufficiently to identify it with either of these species. The thorax in the Panama insect is at least twice as broad as long, finely and closely punctured and pubescent at the sides, nearly smooth and impunctate at the middle of the disc. The scutellum is much broader than long, and closely punctured. The elytra, which are scarcely, if at all, narrowed posteriorly, are distinctly, but not very strongly, punctured anteriorly and very finely so from the middle downwards; the basal third is occupied by a perfectly regular fulvous band of even width, which extends to the lateral and sutural margins and includes the humeral callus; and there is an oval spot close to the apex of each.

E. nigratarsis differs, therefore, from *E. carnifex* by the broad thorax and the black tarsi; and from *E. specularis* by the smaller eyes, the differently punctured red thorax, and the fulvous band of the elytra occupying the entire base. The antennæ are short, and their second and third joints are fulvous; the prosternum is rather broad and of even width.

6. *Euryscopa mexicana*. (Tab. XXXVII. fig. 2.)

Broadly ovate, subcylindrical, black; head and thorax finely pubescent and closely punctured; elytra deeply punctate-striate, with a broad transverse fulvous band at the base, the band not quite extending to the suture nor to the humeral callus.

Length 4 lines.

Head longer than broad, much flattened between the eyes, finely punctured and closely pubescent, the disc piceous; labrum fulvous; antennæ not quite extending to the base of the thorax, black, the third joint fulvous; thorax slightly broader than long, the sides strongly narrowed anteriorly and very much deflexed, the median lobe perfectly straight, the surface very closely and rather finely punctured, clothed at the sides with long whitish pubescence, the disc very sparsely covered; scutellum pubescent and punctured; elytra broad at the base, gradually and slightly narrowed posteriorly, the punctuation very deep anteriorly and fine towards the apex, the fulvous basal band occupying rather more than the anterior third and not quite extending to either the basal or sutural margins, the humeral callus remaining of the black ground-colour and in the shape of a small spot; underside and legs thickly pubescent, black, the tibiæ obscure fulvous, the last abdominal segment with a deep fovea.

Hab. MEXICO, Cuernavaca (*Sallé*).