

A single specimen, apparently a female. *E. mexicana* may be known from its allies by its large size and by the broad fulvous band of the elytra.

7. *Euryscopa cingulata*.

Clythra cingulata, Latr. in Humb. & Bonpl. Obs. Zool. i. p. 208, t. 22. figg. 6, 7 (1811)¹.

Euryscopa cingulata, Lac. Monogr. in Mém. Soc. Liége, v. p. 508².

Megalostomis cingulata, Dej. Cat. 3rd edit. p. 440.

Hab. PANAMA (*Boucard*), Bugaba, Volcan de Chiriqui, Caldera, San Feliz, Tolé, Taboga Island (*Champion*).—COLOMBIA²; ECUADOR, Guayaquil¹.

Many specimens were obtained by Mr. Champion of this apparently not uncommon and handsome species. Examples occur in which the posterior band of the elytra occupies the entire apical portion. Lacordaire, in his excellent description, does not mention that the apical margin of the elytra as well as the suture is narrowly black.

8. *Euryscopa fulvilabris*.

Subcylindrical, black; labrum fulvous; thorax with a longitudinal central ridge, pubescent; elytra strongly punctate-striate, each with a subquadrate red humeral spot enclosing the callus.

Length 2 lines.

Hab. MEXICO, Ventanas in Durango (*Höge*).

Closely allied to *E. scapularis*, Lac., but at once separated by the bright fulvous labrum, the longer antennæ (which extend to the base of the thorax), and the distinctly raised longitudinal line in the centre of the thorax. The scutellum is pubescent, as in *E. scapularis*; but the red humeral spot is quite different in shape, it being nearly square (instead of elongate) and surrounding the callus (this part is always black in *E. scapularis*, and its inner margin is oblique at the base).

A single specimen.

9. *Euryscopa villosa*. (Tab. XXXVII. fig. 3.)

Subcylindrical, black, pubescent; labrum, the basal joints of the antennæ, and the tibiæ fulvous; elytra each with an oblique red spot, finely pubescent, semiregularly and extremely closely punctured, the interstices irregularly rugose.

Length 3 lines.

Head very finely punctured and closely pubescent; labrum fulvous; antennæ dentate within from the third joint, the second, third, and fourth joints fulvous, the rest black, pubescent; thorax about twice as broad as long, closely and finely punctured, clothed with rather short white pubescence, the middle of the disc with a distinct smooth longitudinal line, the median lobe slightly widened and scarcely produced; scutellum with a few punctures and hairs; elytra cylindrical, crowded with larger and smaller punctures, which are irregularly arranged in lines, the interstices sparingly pubescent and slightly convex, the red spot commencing below the humeral callus, gradually narrowing behind, and obliquely extending nearly to the suture; tibiæ fulvous, the femora and tarsi black, pubescent like the underside.

Hab. MEXICO, Chilpancingo and Acapulco in Guerrero (*Höge*).