

sides straight and parallel, slightly converging at the apex, anterior angle subacute, the hinder one acute; anterior margin sinuate on either side; surface impressed with round punctures, which are irregularly scattered over the hinder two thirds of the disc, but are nearly obsolete on the sides and at the apex; flavous; a large subquadrate patch, attached to the basal margin and covering the posterior two thirds of the disc, but not extending to the sides, black; outer limb also narrowly edged with black. Elytra narrowly oblong, parallel, obtusely rounded at the apex, regularly punctate-striate; flavous, a basal patch rufo-testaceous; an entire sutural vitta, broad at its base, but gradually narrowed towards its apex, together with another on each elytron, discoidal, abbreviated at the base but extending downwards nearly to the apex, where it incurves and often joins the sutural vitta, black.

20. *Cephaloleia trivittata*.

Anguste elongata, parallela, modice convexa, nitida; subtus picea, pedibus flavis, piceo-tinctis; supra nigra; thorace subquadrato, subcylindrico, profunde punctato, lateribus rectis, parallelis; elytris fortiter punctato-striatis, flavis, nigro trivittatis, vittis apice confluentibus, duabus externis basi abbreviatis.

Mas. Abdominis segmento anali late emarginato, medio sinuato.

Fem. Abdominis segmento anali late emarginato.

Long. 2 lin.

Hab. GUATEMALA, El Tumbador 2500 feet (*Champion*); PANAMA, Bugaba, Volcan de Chiriqui 2500 to 4000 feet (*Champion*).

Head closely punctured. Antennæ filiform, joints cylindrical, the first slightly thickened, rather longer than the second, about equal in length to the third. Thorax subquadrate; sides straight and parallel, scarcely converging at the extreme apex, anterior angle produced, subacute; disc deeply but not very closely impressed with large round punctures. Scutellum rather longer than broad. Elytra much broader than the thorax, elongate, parallel; apices rounded, slightly sinuate conjointly at the sutural angle; above moderately convex, slightly flattened on the anterior disc; strongly punctate-striate; flavous; with three broad black vittæ, the medial one of which is common, entire, slightly dilated below the scutellum and contracted just before reaching the sutural angle; the others, one on each elytron, commence below the humeral callus, and, after extending the whole length of the disc, incurve to unite with the sutural line.

Much narrower and more elongate than the preceding species.

21. *Cephaloleia semivittata*.

Elongata, nitida, subtus picea; pedibus abdomineque fulvis, hoc piceo tincto; pleuris fortiter punctatis; supra nigra; thorace subquadrato, lateribus rectis; suberebre fortiter punctato; elytris fortiter punctato-striatis, fulvis, vitta suturali communi integra, pone scutellum paullo dilatata, alteraque discoidali a medio ad apicem extensa, apice cum linea suturali connexa, nigris.

Mas. Abdominis segmento anali late transversim emarginato, medio sinuato.

Fem. Abdominis segmento anali apice utrinque sinuato.

Long. 2 lin.

Hab. PANAMA, Bugaba 800 to 1500 feet (*Champion*).