

Head granulose, distinctly punctured. Antennæ less than half the length of the body, robust; basal joint somewhat thickened, scarcely longer than the second, nearly equal in length to the third. Thorax scarcely broader than long; sides straight and parallel, broadly rounded, and abruptly converging at the apex, the anterior angle subacute, the hinder one produced into a small acute tooth; disc subcylindrical, smooth and shining, subremotely impressed with large round punctures. Elytra parallel, rounded at the apex, conjointly subangulate-emarginate at the sutural angle; subnitidous, finely but distinctly punctate-striate, the punctures larger and more strongly impressed on the anterior disc.

38. *Cephaloleia metallescens*.

Ovata, modice convexa, piceo-rufa, nitida; supra obscure cyanea; antennis (basi et apice exceptis) nigris; thorace transverso, sat crebre et fortiter punctato, margine laterali anguste rufo-piceo; elytris sat fortiter punctato-striatis, striis sulcatis, punctis pone medium minus fortiter impressis.

Fem. Abdominis segmento anali rotundato, integro.

Long. $1\frac{2}{3}$ lin.

Hab. GUATEMALA, San Juan in Vera Paz (*Champion*). A single specimen.

Head closely punctured. Antennæ about half the length of the body, robust; first and third joints equal in length, the second slightly shorter; basal joint, together with the apex of the terminal one, rufo-piceous. Thorax nearly twice as broad as long; sides straight, slightly converging from the base to the apex, anterior angle not produced, broadly rounded, hinder angle acute; disc transversely convex, excavated on either side just within the lateral margin, somewhat closely covered with large round punctures; lateral margin narrowly reflexed, rufo-piceous. Scutellum scarcely longer than broad. Elytra broadly oblong-ovate, moderately convex, strongly punctate-striate, the striæ sulcate, the puncturing finer on the hinder disc; interspaces on the lateral margin, and to a less extent on the inner disc, thickened and convex.

Solenispa Weise

39. *Cephaloleia leptomorpha*. (Tab. II fig 4.) - see p. 120

Elongata, angustata, subcylindrica, cuprea, nitida; antennis nigris; pedibus piceo-cupreis; thorace quam lato paullo longiori, lateribus rectis, disco sat remote et tenuiter punctato; elytris parallelis, distincte punctato-striatis.

Mas. Abdominis segmento anali late sed leviter emarginato, medio sinuato.

Fem. Abdominis segmento anali late sed leviter emarginato.

Long. 2 lin.

Hab. PANAMA, Volcan de Chiriqui 8000 feet (*Champion*). (Habits - see p. 120) 8

Vertex and front finely impressed with a double longitudinal groove. Antennæ nearly half the length of the body; basal joint thickened and equal in length to the third, the second joint rather shorter. Thorax distinctly longer than broad; sides straight and parallel, anterior angles rectangular; disc transversely convex, subnitidous, very finely and somewhat sparingly punctured. Elytra broader than the thorax; sides parallel, slightly