

converging near the apex; upper surface slightly flattened along the suture, rather strongly punctate-striate.

The above insect closely resembles some of the species of *Stenispa*.

40. *Cephaloleia tenella*.

Elongata, angustata, parallela, nigra, nitida; dorso æneo vix micans; abdomine piceo, tarsis piceo-fulvis; thorace quam longo vix latiori, lateribus parallelis, disco subcylindrico, sat fortiter punctato; elytris parallelis, apice obtuse rotundatis, sat fortiter punctato-striatis.

Mas. Abdominis segmento anali concavo-emarginato, medio sinuato.

Long. $1\frac{2}{3}$ lin.

Hab. GUATEMALA, Coatepeque 1300 feet (*Champion*).

Vertex and front rather closely punctured. Antennæ filiform, scarcely longer than the head and thorax, slightly thickened towards the apex, the two lower joints equal in length, the basal one not thickened. Thorax very slightly broader than long; sides straight and parallel, abruptly rounded and converging at the apex; anterior margin convex, sinuate on either side; disc transversely convex, rather strongly but not very closely punctured, the puncturing more distant on the medial disc. Elytra rather broader than the thorax, parallel on the sides, obtusely rounded at the apex; disc strongly and regularly punctate-striate, the interspaces impunctate.

41. *Cephaloleia stevensi*. (Tab. I. fig. 20.)

Angustato-elongata, nigra, nitida; pedibus fulvis; thorace sat fortiter punctato, sordide fulvo, vitta discoidali nigra; elytris parallelis, apice obtuse rotundatis, fortiter punctato-striatis, flavis, puncto humerali maculisque rotundatis duabus, communibus, una basi et una vix pone medium positis, nigris.

Long. $1\frac{1}{2}$ lin.

Hab. PANAMA, Bugaba (*Champion*).

Vertex and front closely punctured, the latter concave. Antennæ half the length of the body, filiform, entirely black. Thorax distinctly longer than broad; sides straight and parallel, rounded at the extreme apex, anterior angles not produced, apical margin truncate; upper surface subcylindrical, slightly excavated transversely at the base, strongly and closely punctured; obscure fulvous, a broad longitudinal vitta on the middle disc black, the extreme lateral margin piceous. Scutellum pentangular. Elytra narrowly oblong, parallel, rounded at the apex, strongly punctate-striate, flavous, a small oblong patch on the humeral callus, together with two large common, rotundate spots, one at the base, the other placed just beyond the middle, black.

42. *Cephaloleia sulciceps*.

Sat elongata, subtus fulva, supra sordide rufo-fulva, nitida; antennis (articulis basalibus duobus exceptis) nigris; vertice sat profunde bisulcato; thorace transverso, lateribus parallelis, disco sat remote punctato; elytris parallelis, lateribus ad apicem oblique convergentibus, apice truncatis, dorso convexis, ad latera ante medium sinuatis, regulariter punctato-striatis.