

1. **Prosopodonta limbata.** (Tab. II. fig. 6.)

Prosopodonta limbata, Baly, Cat. Hispidæ, p. 69, t. 6. f. 2¹.

Hab. PANAMA, Volcan de Chiriqui 2500 to 6000 feet (*Champion*).—COLOMBIA¹ (*coll. Baly*, type).

In the specimens from the State of Panama the black colour covers a greater space on the dorsal surface of the elytra than in the type.

2. **Prosopodonta costata.** (Tab. II. fig. 7.)

Balyi Weise 1905.

preacc. by. Waterh. 1879

Oblongo-elongata, subtus nigra, nitida; thorace fulvo-piceo; supra piceo-fulva, antennis nigris; thorace quam longo vix latiori, lateribus leviter rotundatis, basi fere rectis, disco convexo, lævi, basi et ad latera punctato; elytris oblongis, convexis, fortiter punctato-striatis, striis biserialiter dispositis, interspatiis secundo et quarto costatis, costa exteriori [nec non limbo laterali] sordide fulvis.

Long. $3\frac{2}{3}$ lin.

Hab. PANAMA, Volcan de Chiriqui 2000 to 3000 feet (*Champion*).

Head smooth and shining; face armed between the antennæ with an acute tooth. Antennæ rather longer than the head and thorax, distinctly thickened towards the apex, the joints cylindrical. Thorax rather broader than long; sides obtusely rounded, obsolete angulate immediately behind and again just before the middle, straight and slightly diverging at the base, converging from the middle to the apex, apical angle not produced, obtuse; upper surface convex, smooth and shining, impressed on the sides and hinder disc with large deep punctures. Elytra oblong, subparallel, regularly rounded at the apex; strongly punctate-striate, the striæ (nine at the extreme base and eight on the rest of the surface) arranged in double rows; alternate interspaces slightly elevated, the second and fourth from the suture more strongly thickened, costate.

The present insect differs entirely in the coloration of its thorax from any hitherto described species of the genus.

CHEIRISPA.

Cheirispæ, Baly, Cat. Hispidæ, p. 71, t. 1. f. 16.

Four species of this peculiar genus are now known, all from Central America and Colombia.

1. **Cheirispæ dorsata.** (Tab. II. fig. 8.)

Anguste oblonga, nitida, subtus nigra, thorace piceo; supra rufo-picea aut nigro-picea, antennis nigris; facie inter antennas creta elevata, sæpe in dentem curvatum producta, instructa; thorace quam longo latiori, lateribus a basi ad ultra medium fere rectis, parallelis, antice rotundato-angustatis, medio leviter angulatis, disco lævi, ad latera punctato; elytris oblongis, punctato-striatis, striis ad latera et apicem versus