

### 7. *Cephalodonta sallæi*.

*Cephalodonta sallæi*, Baly, Cat. Hispidæ, p. 150<sup>1</sup>.

*Hab.* MEXICO, Cordova<sup>1</sup> (type, coll. Sallé); BRITISH HONDURAS, Rio Hondo (*Blancaneaux*).

### 8. *Cephalodonta pallida*.

Anguste cuneiformis, dorso subdepressa; subtus picea, nitida; pectoris medio, abdominis femorumque basi thoraceque inferiori (hoc utrinque piceo vittato) pallide flavis; supra subopaca, pallide flava; capite lævi, verticis macula, ore antennarumque articulo basali (*cæteri desunt*) nigro-piceis; thorace lateribus obtuse angulatis, rude foveolato-punctato, vitta discoidali nigro-picea; scutello nigro; elytris postice ampliatis, apice obtuse rotundatis, angulo postico dente parvo acuto armato, profunde seriatim punctatis, interspatio quarto valde costato, secundo, sexto et octavo pone medium distincte elevatis; callo humerali lateraliter producto, acuto.

Long. 4 lin.

*Hab.* NICARAGUA, Chontales (*Belt*).

Head smooth; the basal joint of the antennæ short (the rest broken away). Thorax rather broader than long; sides parallel at the base, obtusely angulate in the middle, thence obliquely converging towards the apex, anterior angle subacute; upper surface convex on the sides, broadly flattened on the disc, slightly excavated behind the middle, closely covered with large round punctures; apical margin impunctate. Elytra broader than the thorax; sides increasing in width from the base towards their apex, more abruptly dilated behind the middle, hinder angle armed with a small acute tooth, the apex of which is produced directly backwards; lateral margin entire, the apical margin finely serrulate; upper surface convex on the sides and apex, flattened along the anterior two thirds of the suture; humeral callus laterally produced, its apex acute; each elytron with ten rows of large deep punctures; the fourth interspace strongly costate for its whole length, the second, sixth, and eighth thickened behind the middle of the disc.

The broader thorax and acute apices of the humeral calli will separate this species from *C. javeti*.

### 9. *Cephalodonta gracilenta*. (Tab. II. fig. 21.)

Anguste elongata, parallela, nitida; subtus fulva, thoracis vitta laterali, pectoris lateribus femoribusque piceis, his basi, tibiis tarsisque pallide rufo-fulvis, abdomine piceo tincto; supra flava, antennis nigro-piceis aut nigris, apice sordide rufo-piceis; thorace crebre foveolato-punctato, medio nigro-piceo vittato, lateribus vix pone medium angulatis; scutello nigro-piceo; elytris elongatis, parallelis, medio leviter constrictis, apice obtuse rotundatis, angulo postico spina acuta armato, fortiter seriatim punctatis, utrisque vitta humerali, fere ad medium extensa, anguloque postico nigro-piceis.

*Mas.* Tibiis anticis intus prope apicem spina aut dente brevi armatis.

*Fem.* Tibiis anticis intus prope apicem tuberculatis.

Long. 3-3½ lin.

*Hab.* PANAMA, Bugaba 800 to 1500 feet (*Champion*).

Vertex and front shining, impunctate, the latter impressed with an oblong fovea; the