

Hab. GUATEMALA (type in coll. Baly ex coll. Laferté); PANAMA, David, Taboga Island (*Champion*).—COLOMBIA¹.

3. *Charistena trilineata*.

Charistena trilineata, Baly, Trans. Ent. Soc. 1864, p. 255¹.

Hab. MEXICO¹ (types in coll. Chevr. and Baly).

4. *Charistena pilatei*.

Charistena pilatei, Baly, Trans. Ent. Soc. 1864, p. 254¹.

Hab. MEXICO, Teapa¹ (coll. Baly, type); GUATEMALA, Chiacam in Vera Paz (*Champion*).

5. *Charistena championi*.

Angustata, subcylindrica; subtus cum capite nigra, pedibus flavis, antennis basi tarsisque pallide-piceis; supra flava, scutello nigro; thorace subcylindrico, profunde et crebre punctato, lateribus plaga margineque apicali nigris; elytris parallelis, apice rotundatis, minute serrulatis; utrisque octo-seriato-punctatis, interspatiis secundo, quarto sextoque, nec non sutura, modice costatis; maculis communibus duabus, una ad basin altera vix ante apicem positis, vittaque brevi submarginali baseos, nigris, æneo tinctis.

Var. A. Elytrorum macula subapicali obsoleta.

Long. $1\frac{1}{3}$ lin.

Hab. GUATEMALA, Zapote, Tamahu in Vera Paz (*Champion*).

Head tinged with æneous; vertex and front impressed with a number of longitudinal grooves. Antennæ slightly longer than the head and thorax, robust, thickened towards the apex; joints cylindrical, the five upper ones forming an ill-defined elongate club; six lower joints rufo-piceous, the rest black. Thorax subcylindrical, rather broader than long; sides obtusely angulate, straight and parallel behind the middle, obliquely converging anteriorly; upper surface deeply and closely punctured; a patch on the lateral margin, together with the extreme apex, black. Elytra parallel, regularly rounded at the apex, faintly sinuate at the sutural angle, distinctly serrulate; each elytron with eight, at the extreme base with nine, rows of large deep punctures; the second, fourth, and sixth interspaces moderately costate, the suture also thickened.

6. *Charistena funesta*.

Angustata, subcylindrica, nigra, nitida, dorso æneo tincta; thorace profunde foveolato-punctato, disco anteriori fere impunctato; elytris parallelis, tricostatis, interspatiis profunde biseriato-punctatis.

Long. 2 lin.

Hab. MEXICO, Orizaba (*Sallé*); GUATEMALA, San Gerónimo (*Champion*).

Head very finely granulose, front impressed between the eyes with a faint elongate fovea. Antennæ equal in length to the head and thorax; five terminal joints thickened, forming a narrow elongate club. Thorax longer than broad; sides parallel, very slightly angulate; subcylindrical, excavated at the base just in front of the scutellum,