

covered with large, round, foveolate punctures; medial line with a short longitudinal groove. Elytra subparallel, slightly increasing in width towards the apex; lateral margin finely serrulate; the apical margin dilated, regularly rounded, broadly quadrate-emarginate at the sutural angle, more coarsely serrulate than the sides; moderately convex, slightly flattened along the suture; each elytron with eight longitudinal rows of large, deeply impressed punctures; the second, fourth, and sixth interspaces, together with the suture, costate.

2. *Apices of the elytra entire, or only obsoletely sinuate at the sutural angle.*

3. *Chalepus semilimbatus.*

Elongatus, angustatus, parallelus, flavus, nitidus; oculorum orbitis, antennis pedibusque (femoribus basi exceptis) nigris; thorace subcylindrico, fortiter punctato, lateribus, pone medium fere rectis, ante medium constrictis; elytris parallelis, apice rotundatis, minute sed distincte serrulatis; utrisque octo-seriato-punctatis, stria sexta medio obsoleta; interspatiis alternis costatis; lateribus anguste nigro-æneo marginatis.

Long. $2\frac{1}{2}$ lin.

Hab. MEXICO, Vera Cruz (*Sallé*).

Vertex and front impressed with a deep longitudinal fovea; interocular space moderately produced. Antennæ about one third the length of the body, slightly but distinctly thickened towards the apex; joints cylindrical. Thorax transverse; sides rounded, nearly straight and parallel behind the middle, thence rounded and converging towards the apex, deeply sinuate behind the anterior angle, the latter produced laterally into a short, acute tooth; transversely convex, rather deeply excavated in front of the scutellum, deeply and rather closely punctured. Elytra subelongate, parallel, obtusely rounded at the apex, finely serrulate; each elytron with eight rows of deep punctures, the sixth row obsolete on the middle disc; second and fourth interspaces broadly, the sixth less strongly, costate; lateral margin narrowly edged with blackish æneous.

4. *Chalepus suturalis.*

Anguste oblongus, subtus niger, nitidus; supra rufo-fulvus, minus nitidus; capite nitido elytrorumque linea suturali nigro-æneis, scutello antennisque nigris; thorace transverso, rude et crebre punctato, lateribus angulatis; elytris oblongis, parallelis, apice rotundatis, utrisque octo-seriato-punctatis, interspatiis alternis vix elevatis.

Long. 2 lin.

Hab. GUATEMALA, Dueñas (*Champion*).

Vertex smooth, front trisulcate; interocular space slightly produced. Antennæ nearly half the length of the body, robust, slightly thickened towards the apex; joints cylindrical. Thorax transverse; sides angulate, straight and parallel from the base to the middle, thence obliquely converging to the apex, the anterior angle acute; disc transversely convex, slightly excavated behind the middle, coarsely rugose-punctate. Elytra narrowly oblong; sides parallel, obtusely rounded at the apex; lateral margin