

Head smooth, impunctate; front impressed with three short longitudinal grooves; interocular space moderately produced. Antennæ nearly half the length of the body, filiform, rather slender, slightly thickened towards the apex; joints cylindrical, first and second short, equal, slightly thickened. Thorax rather broader than long, subconical; sides converging from base to apex; transversely convex, transversely depressed on the hinder disc, closely punctured; the hinder margin elevated. Elytra broader than the thorax; sides parallel, very slightly enlarged towards the posterior angle; apex obtusely rounded, outer margin nearly entire, obsoletely serrulate behind the middle and at the apex; each elytron with eight, at the extreme base with nine, rows of large, deep punctures; second, fourth, and sixth interspaces moderately costate.

Resembles in coloration *C. congener*, but half the size, and rather less parallel in form, the elytra with an additional row of punctures at the base.

11. *Chalepus tibialis*.

Angustatus, elongatus, subcylindricus; subtus niger, nitidus; femoribus basi, anticis fere totis, tibiis anticis basi, posticis fere totis, pectoris medio abdominisque disco, flavis; supra flavus, subnitidus, scutello capiteque nigris, facie inter oculos flava; thorace transverso, crebre punctato, lateribus vittaque lata discoidali nigris; elytris parallelis, octo-basi nono-seriato-punctatis, tricostatis, linea laterali baseos vittaque suturali nigris; tibiis anticis intus ante apicem spina brevi armatis.

Long. 2 lin.

Hab. PANAMA, Volcan de Chiriqui 2500 to 4000 feet (*Champion*).

Head smooth, slightly produced between the eyes. Antennæ nearly half the length of the body, slightly thickened towards the apex; joints cylindrical, the two lower ones short, nearly equal, the third distinctly longer. Thorax broader than long; sides obsoletely angulate, straight and parallel from the base to the middle, thence converging and slightly sinuate to the apex; anterior angle obliquely produced into a distinct subacute tooth; transversely convex, faintly excavated transversely behind the middle, coarsely and closely foveolate-punctate, the punctures rather more distant on the medial line; the lateral margin, together with a broad discoidal vitta, black, the apical margin narrowly edged with piceous. Elytra rather broader than the thorax; sides parallel, very faintly constricted in the middle, obsoletely serrulate; apex regularly rounded, its serratures rather more distinct; each elytron with eight, at the extreme base with nine, rows of punctures; the second, fourth, and sixth interspaces costate.

12. *Chalepus jansonii*.

Subelongatus, subtus cum capite niger, nitidus; pedibus maculaque frontali fulvis; supra fulvus, minus nitidus, scutello nigro; thorace subcylindrico, rude et crebre punctato; elytris anguste oblongis, parallelis, apice rotundatis, margine serrulatis; utrisque octo-basi nono-seriato-punctatis, interspatiis secundo, quarto sextoque nec non sutura modice costatis.

Long. $1\frac{1}{2}$ lin.

Hab. NICARAGUA, Chontales (*Janson*).

Vertex finely granulose, front longitudinally grooved; interocular space moderately