

Antennæ robust, obsolete thickened towards the apex, rather longer than the head and thorax; joints cylindrical, the first two short, equal, the third scarcely longer than the second. Thorax broader than long, sides converging from base to apex, subangulate-rotundate; disc transversely convex, transversely excavated behind the middle, coarsely and closely punctured. Elytra broader than the thorax; sides very slightly dilated, the apices regularly rounded, outer margin distinctly serrulate; each elytron at the extreme base and on the hinder disc with ten, on the anterior disc with nine, rows of large, deeply impressed punctures; second and fourth interspaces rather strongly and equally costate, the eighth interspace less strongly elevated, the sixth interspace thickened at its apex, the suture also thickened.

28. *Chalepus pascoei*.

Subelongatus; subtus niger, nitidus; pectoris medio, femoribusque basi fulvis; supra flavo-fulvus, minus nitidus; antennis, scutello, nec non elytrorum vitta suturali baseos nigris; thorace transverso, lateribus angulatis; disco crebre, minus fortiter punctato; lateribus angustis vittaque discoidali, hac sæpe basi et apice abbreviata, nigris; elytris fere parallelis, apice rotundatis, margine minute serrulato; utrisque decem-paullo ante medium nono-seriato-punctatis; interspatiis secundo et quarto æqualiter, octavo minus distincte, costatis.

Long. $2\frac{1}{2}$ lin.

Hab. GUATEMALA, San Gerónimo (*Champion*).

Head smooth, impunctate; vertex impressed with a shallow longitudinal groove; interocular space moderately produced. Antennæ robust, rather longer than the head and thorax, slightly thickened towards the apex; joints cylindrical. Thorax transverse; sides angulate, converging from base to apex; disc transversely convex, transversely excavated behind the middle, closely and coarsely, but not very deeply punctured. Elytra broader than the thorax, subparallel, regularly rounded at the apex, the outer margin minutely serrulate; each elytron with ten, just before its middle with eight or nine, rows of punctures; the second and fourth interspaces costate, the eighth less strongly elevated.

29. *Chalepus amabilis*.

Anguste elongatus, parallelus, flavo-fulvus; subtus læte fulvus, nitidus; pectoris lateribus abdominisque apice nigro-piceis, pedibus nigris, femoribus (apice excepto) piceo-fulvis; supra opacus, capite (vertice excepto) nigro; thorace conico, lateribus fere rectis, medio obsolete angulatis, anguste nigro marginatis; disco subcylindrico, pone medium transversim excavato, crebre rugoso-punctato; elytris parallelis, apice rotundatis, lateribus minute serrulatis, margine apicali paullo dilatato, acute denticulato; utrisque decem-prope medium nono-seriato-punctatis; interspatiis secundo et quarto æqualiter, octavo minus fortiter, costatis.

Long. $3\frac{1}{4}$ lin.

Hab. MEXICO, Tehuantepec (*Sallé*). A single specimen.

Face strongly produced between the eyes; front deeply trisulcate. Antennæ moderately robust, filiform, slightly thickened towards the apex; joints cylindrical, the third