

Head moderately produced between the eyes; front longitudinally sulcate on the medial line, irregularly sculptured on either side. Antennæ half the length of the body, slightly thickened towards the apex; the two lower joints ovate, equal, the third rather longer. Thorax longer than broad; sides rounded, nearly straight and parallel from the base to the middle, sinuate just behind the latter, constricted in front, the anterior angle produced obliquely into a subacute tooth; upper surface transversely convex, slightly excavated on the hinder disc, closely foveolate-punctate; the lateral margin, together with a broad discoidal vitta, black. Elytra broader than the thorax; sides parallel, obsoletely constricted in the middle, the apex regularly rounded; lateral margin obsoletely, the apical one more distinctly, serrulate; each elytron with ten rows of punctures, the second, fourth, sixth, and eighth interspaces costate, the sixth less strongly raised, its margin serrulate.

#### 47. *Chalepus marginatus*.

Elongatus, subtus niger, nitidus; thorace, lateribus exceptis, pectoris medio femoribusque anticis quatuor basi, fulvis; supra subnitidus, fulvus; scutello capiteque nigris, hoc macula frontali sordide fulva; thorace lateribus rotundatis, antice constrictis; rude rugoso-punctato, vitta discoidali lateribusque nigris; elytris parallelis, apice rotundatis, tenuiter serrulatis; utrisque decem-seriato-punctatis; interspatiis secundo quartoque modice, octavo obsolete, costatis; limbo laterali anguste nigro, sutura ante medium anguste nigro-picea.

Long. 2 lin.

*Hab.* MEXICO, Cordova (*Sallé*). A single specimen.

Front moderately produced between the eyes, longitudinally sulcate. Antennæ nearly half the length of the body, subfiliform, slightly thickened towards the apex; joints cylindrical. Thorax transverse; sides rounded, obliquely diverging from the base to the middle, deeply sinuate behind the apex, the apical angle armed laterally with an acute tooth; disc coarsely and strongly punctured, transversely excavated behind the middle. Elytra parallel, regularly rounded at the apex; lateral margin obsoletely, the apical margin finely, serrulate; each elytron with ten regular rows of punctures, the second and fourth interspaces moderately costate, the eighth obsoletely thickened.

#### 48. *Chalepus lateralis*.

Elongatus, parallelus; subtus niger, nitidus; thoracis lateribus, abdominis disco, femoribus basi tibiæque annulo lato, fulvis; supra subopacus, fulvus, scutello capiteque nigris; thorace transverso, lateribus rotundatis, pone medium fere rectis, ante medium sat abrupte constrictis; disco transversim convexo, pone medium transversim excavato, rude et crebre punctato; lateribus maculaque vix pone medium posita, nigris; elytris parallelis, apice rotundatis, ad apicem minute serrulatis; utrisque decem-seriato-punctatis, sutura nec non interspatiis secundo et quarto modice costatis, interspatio octavo minus distincte elevato; utrisque linea laterali, a basi ad longe pone medium extensa, nigra ornatis.

Long. 2 lin.

*Hab.* GUATEMALA, Zapote, Capetillo, Panima in Vera Paz (*Champion*).

Vertex smooth, impunctate; front irregularly sulcate, subrugose; interocular space