

12. *Physonota separata*.

Physonota separata, Boh. Monogr. Cassid. ii. p. 205¹, and iv. p. 249²; Cat. Col. Ins. Brit. Mus. ix. p. 99³.

Cassida nigropunctata, Sturm, in litt.⁴

Deloyala pallida, Sturm, in litt.⁵

Hab. MEXICO²⁻⁵ (*Chevrolat*¹; *Sallé, ex coll. Sturm*), Cordova (*Sallé*), Cuernavaca (*Höge*).

I have seen four specimens of this insect, including one, no doubt the type, from the Chevrolat collection in the British Museum. *P. separata* only differs from *P. vitticollis* in having much fewer punctures on the disc of the elytra, these being widely separate one from another.

13. *Physonota turgida*. (Tab. VIII. fig. 20.)

Physonota turgida, Boh. Monogr. Cassid. ii. p. 197¹, and iv. p. 246²; Cat. Col. Ins. Brit. Mus. ix. p. 99³.

Physonota bicolor, Sturm, in litt.

Hab. MEXICO¹⁻³ (*Sallé, ex coll. Sturm*), Oaxaca (*Höge*).

There are two examples of this peculiar species in the Sallé collection, without nearer indication of locality. The type from the Stockholm Museum is figured. A rather narrow, oval species, with coarse, subserially arranged punctures on the elytra, the margins also coarsely punctured. The single specimen from Oaxaca is much smaller than the type, and has the dorsum of the elytra pitchy-brown, probably from discoloration.

14. *Physonota mexicana*. (Tab. VIII. fig. 21.)

Physonota mexicana, Boh. Monogr. Cassid. ii. p. 206¹, and iv. p. 249²; Cat. Col. Ins. Brit. Mus. ix. p. 100³.

Hab. MEXICO¹⁻³, Guanajuato (*Sallé*), Tula and Pachuca in Hidalgo (*Höge*).

Sent in plenty by Höge from the State of Hidalgo. *P. mexicana* is closely allied to *P. ovata*, Boh., but may easily be known from it by the blackish arcuate or >-shaped mark on either side of the disc of the prothorax near the base, midway between which is a short oblong streak; the elytra are rather coarsely, irregularly, seriate punctate, the margins also coarsely punctured.

15. *Physonota flavago*. (Tab. VIII. fig. 22.)

Physonota flavago, Boh. Monogr. Cassid. ii. p. 210¹, and iv. p. 249²; Cat. Col. Ins. Brit. Mus. ix. p. 100³.

Hab. MEXICO¹⁻³, Ciudad in Durango 8100 feet (*Forrer*).

We figure the type from the Stockholm Museum. Very closely allied to *P. picticollis*, Boh., but differing from it in the immaculate prothorax, and in having the elytra more coarsely and more closely punctured. The two specimens (♂ ♀) received