

narrowly edged with sanguineous, the punctures on the outer part of the disc coarser and more remote from each other. A specimen from Rio Hondo is figured.

17. **Coptocycla atalanta.** (Tab. X. fig. 14.)

Coptocycla atalanta, Boh. Monogr. Cassid. iv. p. 416¹.

Hab. MEXICO, San Andres Tuxtla and Playa Vicente, both in Vera Cruz (*Sallé*¹), Misantla (*Höge*), Teapa in Tabasco (*H. H. Smith*); BRITISH HONDURAS, R. Hondo (*Blancaneaux*); GUATEMALA, Senahu in Vera Paz (*Champion*).

Apparently confined to the Atlantic slope. This species again is closely allied to *C. opulenta*, Boh.; but it is smaller, with the annulus narrower (sanguineous, bordered externally with black), and the suture narrowly sanguineous nearly to the apex, the sutural stripe not ovately dilated below the base. The third joint of the antennæ is elongate. We figure a specimen from Misantla.

18. **Coptocycla extensa.** (Tab. X. fig. 15, ♀.)

Coptocycla extensa, Boh. Monogr. Cassid. iii. p. 261¹, and iv. p. 435²; Cat. Col. Ins. Brit. Mus. ix. p. 175³.

Var. β . The elytra with the disc testaceous or flavo-testaceous.

Hab. MEXICO¹⁻³ (*coll. Oberthür*), Teapa in Tabasco (*H. H. Smith*), Temax in North Yucatan (*Gaumer*), Yucatan (*coll. Baly*).

Sent in plenty from Yucatan, about one-third of the specimens belonging to the var. *a* of Boheman—that is to say, with the disc of the elytra fusco-testaceous or castaneous, without distinct yellow markings.

19. **Coptocycla zetterstedti.** (Tab. X. fig. 16, ♂.)

Coptocycla zetterstedti, Boh. Monogr. Cassid. iii. p. 303¹, and iv. p. 443²; Cat. Col. Ins. Brit. Mus. ix. p. 180³.

Hab. MEXICO, Vera Cruz, Panistlahuaca (*Sallé*), Cordova (*Sallé*, *Höge*), Acapulco, Misantla, Tapachula (*Höge*).

Not uncommon at Cordova and elsewhere in the Mexican State of Vera Cruz. The antennæ are elongate and slender, the third joint nearly twice as long as the second. The single specimen (immature) received from Tapachula has the prothorax almost immaculate. Our figure is taken from a male example from Cordova.

20. **Coptocycla decussata.**

Coptocycla decussata, Boh. Monogr. Cassid. iii. p. 305¹, and iv. p. 443²; Cat. Col. Ins. Brit. Mus. ix. p. 180³.

“Subrotundata, modice convexa, flava, nitida; antennis articulis duobus ultimis nigro-fuscis; prothorace subtilissime, subremote punctulato, antice late rotundato, pone medium maculis tribus parvis, transverse