

40. **Coptocycla semiatrata.** (Tab. XI. fig. 8.)

Coptocycla semiatrata, Boh. Monogr. Cassid. iv. p. 428¹.

Hab. MEXICO, Cordova (*Sallé*¹), Atoyac in Vera Cruz, Teapa in Tabasco (*H. H. Smith*); GUATEMALA, Las Mercedes, Cerro Zunil (*Champion*).

This insect is exceedingly like the similarly-coloured *Ctenochira plicata* (Boh.), but, apart from the non-pectinate claws, it may be distinguished by the smoother and less coarsely punctured elytra, the disc of the latter not deeply foveate on either side of the suture a little below the base. The third joint of the antennæ is not longer than the second. A specimen from Teapa is figured.

41. **Coptocycla crucipennis.** (Tab. XI. fig. 9.)

Coptocycla crucipennis, Boh. Monogr. Cassid. iii. p. 307¹, and iv. p. 444²; Cat. Col. Ins. Brit. Mus. ix. p. 180³.

Hab. MEXICO¹⁻³, San Andres Tuxtla, Playa Vicente (*Sallé*), Cordova, Misantla, Oaxaca, Tapachula (*Höge*), Atoyac, Teapa (*H. H. Smith*); BRITISH HONDURAS, R. Hondo (*Blancaneaux*); GUATEMALA, Yzabal (*Sallé*), Panima, Panzos, San Juan, Senahu, Tactic, and San Gerónimo in Vera Paz, Capetillo, Zapote (*Champion*); HONDURAS (*Sallé*); NICARAGUA, Granada (*Sallé*); COSTA RICA (*Rogers, Van Patten*).

Not uncommon in Mexico and Guatemala. The antennæ usually have the apical joint infusate or black, and sometimes the penultimate joint also; in some specimens, however, only the extreme tip of the apical joint is infusate; the third joint is short, very little longer than the second. The elytra are rather feebly emarginate at the base. A specimen from Misantla is figured.

42. **Coptocycla dohrni.** (Tab. XI. fig. 10.)

Coptocycla dohrni, Boh. Monogr. Cassid. iii. p. 323¹, and iv. p. 447²; Cat. Col. Ins. Brit. Mus. ix. p. 182³.

Hab. MEXICO¹⁻³, Matamoros Izucar, Orizaba, Vera Cruz (*Sallé*), Cordova (*Sallé, Höge*), Misantla, Jalapa (*Höge*), Atoyac in Vera Cruz (*H. H. Smith*).

This insect only differs from *C. crucipennis* in having the anterior ramus of the broad C-shaped mark on the outer part of each elytron obliterated (not extending across the margins to the humeri, as in *C. crucipennis*), and the other elytral markings reduced in size. One specimen from Cordova is almost intermediate. An example from Jalapa is figured.

43. **Coptocycla distorta.** (Tab. XI. fig. 11.)

Coptocycla distorta, Boh. Monogr. Cassid. iii. p. 324¹, and iv. p. 447²; Cat. Col. Ins. Brit. Mus. ix. p. 182³.

Hab. MEXICO¹⁻³, Yolotepec in Oaxaca (*Sallé*).