

*C. incincta* has comparatively short, subfiliform antennæ, with joint 3 shorter and thinner than 2, 4 fully twice as long as 3, 5-10 slightly decreasing in length. The deep punctures on the elytra are rather widely separated from each other, especially on the rounded black patch enclosed by the yellow annulus.

46. **Coptocycla circumducta.** (Tab. XI. figg. 14; 14 *a*, antenna.)

*Coptocycla circumducta*, Boh. Monogr. Cassid. iii. p. 161<sup>1</sup>, and iv. p. 406<sup>2</sup>; Cat. Col. Ins. Brit. Mus. ix. p. 165<sup>3</sup>.

*Hab.* PANAMA, Volcan de Chiriqui (*Champion*).—BRAZIL<sup>1</sup>, Rio Grande<sup>2 3</sup>.

Three male examples from Chiriqui, agreeing with others from Brazil. No mention is made by Boheman of the sexual differences, the males having, as in some of the allied forms, the elytra very obtuse and almost separately rounded at the apex. The antennæ are moderately elongate and subfiliform, with joint 3 much shorter and thinner than 2, 4 twice as long as 3, 5 nearly twice as long as 4, 6-10 subequal in length, 11 very elongate, black at the tip.

47. **Coptocycla aurofasciata.** (Tab. XI. figg. 15, ♂; 16, var. *fåhræi*, ♀.)

*Cassida aurofasciata*, Er. in Wieg. Archiv f. Naturg. xiii. 1, p. 155<sup>1</sup>.

*Coptocycla aurofasciata*, Boh. Monogr. Cassid. iii. p. 224<sup>2</sup>, and iv. p. 429<sup>3</sup>; Cat. Col. Ins. Brit. Mus. ix. p. 171<sup>4</sup>.

*Hab.* PANAMA (*Boucard*), Bugaba (*Champion*).—SOUTH AMERICA, Cayenne<sup>2-4</sup>, Amazons, Brazil<sup>2-4</sup>, Peru<sup>1-4</sup>.

Var. *α*. The transverse fascia on the elytra broader; antennæ with joints 2-10 infusate beneath.

*Hab.* NICARAGUA, Chontales (*Belt*).

Var. *β*. The elytral fascia extending forwards along the first and second interstices to the base, leaving a large rounded patch on each elytron, and nearly the apical half, black; antennæ with joints 4-6 and 11 infusate at the tip beneath.

*Coptocycla fåhræi*, Boh. Monogr. Cassid. iii. p. 229<sup>5</sup>, and iv. p. 429<sup>6</sup>; Cat. Col. Ins. Brit. Mus. ix. p. 172<sup>7</sup>.

*Hab.* MEXICO, Teapa (*coll. Baly*); GUATEMALA, Panzos in Vera Paz (*Champion, Conradt*); HONDURAS<sup>5-7</sup>.

*C. fåhræi*, the type (♀) of which we figure (fig. 16), is evidently only a variety of the widely distributed *C. aurofasciata*. The males, as in the closely-allied *C. leprieuri*, Boh., differ from the females in having the elytra broadly subtruncate at the apex. The antennæ are elongate and subfiliform, with the third joint (as in *C. leprieuri*) distinctly shorter and thinner than the second, the fourth twice as long as the third, the fifth more than one-half longer than the fourth, joints 5-10 being elongate and subequal in length.