

## 2. *Crotchia angustula*.

Angustata, ænea, nitida; capite et prothorace minute parcius punctatis, hoc quadrato, angulis anticis subcallosis et parum prominulis vel reflexis; elytris fortius punctato-striatis; antennis pedibusque nigris, illis basi piceis. Long. 6-7 millim. ♂ ♀.

Mas tarsis anticis latis, villosis; abdominis segmentis tribus intermediis puncto duplici setigero munitis.

Femina tarsis anticis articulo basali parvo.

*Hab.* PANAMA, Bugaba (*Champion*).

With the same general structure as *C. proxima* this species differs from it in being much smaller and narrower; and in the striæ of the elytra having fewer, but much deeper and larger, punctures. The tarsi also differ more in the sexes. The antennæ are more or less picescent, excepting the club. The body beneath in both this and the preceding species is woolly black and shining, the apical ventral segment alone being somewhat pubescent; the ventral segments are vaguely impressed on the sides, but the setigerous spot in the males is placed a little on each side of the middle. About a dozen examples were secured by Mr. Champion.

## 3. *Crotchia veræpaci*s.

*C. parallelæ* (Fowler) affinis; elongata, angusta, picea, nitida; capite prothoraceque parcius sat distincte punctatis, hoc quadrato, basi leviter sinuato; elytris fortius punctato-striatis; metasterno punctato; antennis clava nigra. Long. 5 millim. ♀.

*Hab.* GUATEMALA, El Jicaro in Vera Paz (*Champion*).

This insect is so very nearly allied to a species in my collection, from the Amazons, described by Mr. Fowler as *C. parallela*, that I have some hesitation in treating it as distinct. Unfortunately there is only a single female specimen; but having cleaned and carefully compared the types of both, I am able to state that *C. veræpaci*s differs in having the metasternum distinctly and sparingly punctate, the head and thorax rather less thickly punctured, and the elytra have the punctures larger. I do not at present detect any other points by which I can distinguish *C. veræpaci*s from the female of *C. parallela*. It is about the size of a small *C. angustula*, but is even narrower and more parallel than that species.

## 4. *Crotchia curvipes*.

Picea vel nigro-ænea, nitida; capite parcius sat fortiter, prothorace parce subtiliter, punctatis, hoc subquadrato; elytris leviter punctato-striatis; antennis rufo-piceis, clava nigra. Long. 6 millim. ♂ ♀.

Mas abdominis segmentis intermediis puncto duplici setigero munitis; tibiis intermediis curvatis.

*Hab.* PANAMA, Bugaba (*Champion*).

The male characters of this species will distinguish it at once from *C. hondurana*; it is also larger than that insect, which it otherwise resembles in the punctuation. The front angles of the thorax are a little more decidedly callous and turned outwards; the base of the thorax has a few large scattered punctures in the depressed portion, the punctiform basal sulcus deep and evident; and the legs are more robust and more evidently punctured.