

parvulis in margine antica, limbo laterali, elytris macula magna communi tribusque aliis (una subhumerali, duabus paullo post medium transverse dispositis), et sutura nigris. Long. 6 millim.

Hab. PANAMA, Bugaba (*Champion*).

Very nearly allied to *I. septemsignatus*, but more obsoletely punctured and with the markings different. The head has no spot; the thorax has two small spots upon the front margin, and no basal marks of any kind. The two spots on the middle of the elytra do not represent the one spot in *I. septemsignatus*; they are placed further back. The suture is very narrowly black, and that colour vanishes before the apex. The underside and legs are pale, but possibly the unique specimen is not fully matured.

One specimen.

13. *Ischyrus undulatus*. (Tab. III. fig. 3.)

Oblongo-ovatus, flavo-ferrugineus, subtus subtiliter punctatus; ore, antennarum clava, prothoracis punctis quatuor (duobus anterioribus approximatis, duobus posterioribus magis distantibus), elytris macula magna scutellari communi, puncto humerali et subapicali fasciaque mediana, valde attenuata, marginem haud attingente, sutura pedibusque nigris. Long. 7 millim.

Hab. PANAMA, Bugaba (*Champion*).

Head red, finely and thickly punctured, the front edge blackish; antennæ pitchy, their third joint and the club black; thorax less densely and more deeply punctured than the head, the four spots arranged in a trapezium, the two front ones being nearest, and not touching the front margin. The punctured striæ of the elytra are very faintly impressed and obliterated at the sides and apex.

This and *I. nitidior*, Crotch, are the only species of *Ischyrus* at present known to me with the spots arranged on the thorax in this way. Allied to *I. nitidior*, Crotch, but larger and differently punctured. A single specimen.

14. *Ischyrus pictus*. (Tab. III. fig. 4.)

Oblongo-ovatus, crebre sat fortiter punctatus, flavus; vertice, antennis, prothoracis punctis duobus discoidalibus et tribus basalibus, scutello, elytris macula scutellari communi subquadrata, gutta oblonga juxta suturam pone medium, punctisque tribus (una humerali, una mediana prope marginem et una subapicali), sutura, limbo tenui pedibusque nigris; pectore picescente. Long. 6 millim.

Hab. GUATEMALA, San Juan in Vera Paz (*Champion*).

Rather broadly ovate and convex; the puncturing moderately strong, the striæ evanescent at the sides and apex of the elytra, but fine interstitial puncturing may be seen. The mouth and palpi are pitchy; the head is yellow above, but the base and the whole margin of the epistome are blackish. The thorax is about twice as wide as long, the sides not much rounded, narrowing in front; the three basal spots are connected along the base, the two outside ones being somewhat triangular. The tarsi are (as usual) pitchy-red. The markings of the elytra need not be re-enumerated, as the figure gives a very good representation of them. A single specimen.