

15. **Ischyrus insolens.** (Tab. II. fig. 20.)

*Ischyrus insolens*, Crotch, Cist. Ent. i. p. 429<sup>1</sup>.

*Hab.* MEXICO<sup>1</sup>, Toxpam (*Sallé*).

This is one of the largest of the true *Ischyri*, and will be recognized by a reference to our Plate, as the markings are peculiar. It has, so far as I know, only been met with by M. Sallé.

16. **Ischyrus ehippiatus.**

Oblongo-ovatus, subtus subtiliter, supra fortius parce punctatus; niger, capite prothoraceque rufis, hoc vitta lata et basi nigris, elytris rufis plagia late marginem haud attingente, sutura limboque tenui nigris; tarsis rufis. Long.  $5\frac{1}{2}$  millim.

*Hab.* PANAMA, Bugaba, Caldera in Chiriqui (*Champion*).

Allied to *I. auriculatus*, Lac., from Cayenne, in the mode in which the thorax is marked with black; the head is rufous, impressed in front, almost coarsely punctured; the antennæ are black; the thorax is less thickly but strongly punctured. Elytra narrowing behind, punctate-striate; in addition to the markings mentioned in the diagnosis, the extreme limb is black; the suture is black throughout; the black discoidal patch is indented posteriorly at the suture (and looks as if it would divide into a single vitta on each elytron), and again indented between the fourth and fifth striæ, both in front and behind. A single specimen from each locality.

17. **Ischyrus chacojæ.** (Tab. III. fig. 5.)

Sanguineo-rufus, capite et corpore subtus nigrescentibus, sat fortiter punctatus; antennis, pedibus, pronoti punctis duobus, sutura fasciisque duabus extus abbreviatis (una basali juxta suturam latiore, altera pone medium), nigris. Long.  $5\frac{1}{2}$  millim.

*Hab.* GUATEMALA, Chacoj in Vera Paz (*Champion*).

The head in this species is black, obscurely dark red in the middle; the thorax is dark blood-red, with two black dots placed rather near together in the middle, but nearer the front margin than to the base. The elytra are black at the base, but not externally to the callus, and more widely so to the third stria; the suture is black throughout; and there is an abbreviated fascia beyond the middle somewhat constricted near the suture; the striæ have large and deep punctures at the base, and they are also distinct beyond the middle, but the apex is nearly smooth. The legs are black, the claws and claw-joint red. A single specimen.

18. **Ischyrus quinquepunctatus.** (Tab. III. fig. 6.)

Niger, prothorace rufo, punctis duobus discoidalibus et tribus basalibus nigris; elytris sat fortiter punctato-striatis, fasciis duabus ad marginem latioribus et fere conjunctis tarsisque rufis. Long. 5 millim.

*Hab.* PANAMA, Bugaba (*Champion*).