

Crotch appears to have known this species, but it is not represented in the Cambridge collection. There are, however, in our collection several specimens which agree very well with the description, and which I accordingly refer to it, besides some others sent by Morrison. Fabricius<sup>1</sup> gives as locality "America meridionalis," and our insect has certainly a wide range in Central America. The specimens from Sonora differ from the others in having the lateral spot on the elytra elongate, reaching almost to the apical one. *H. sexverrucata* is not included by Crotch among the Hyperaspides of the United States.

We figure a specimen from Chilpancingo.

17. **Hyperaspis kunzii.** (Tab. XI. fig. 5, ♂.)

*H. gorhami* Kor. k

*Hyperaspis kunzii*, Muls. Spec. Col. Trim. sécur. p. 672<sup>1</sup>; Crotch, Rev. Coccin. p. 237<sup>2</sup>.

*Hab.* MEXICO, Guanajuato (*Sallé*).

The locality for this species was unknown to Mulsant<sup>1</sup> and Crotch<sup>2</sup>. There are two examples of it from Mexico in the *Sallé* collection.

vi. *Elytra with two yellow spots.*

18. **Hyperaspis deyrollii.**

*Hyperaspis deyrollii*, Crotch, Rev. Coccin. p. 229<sup>1</sup>.

*Hab.* MEXICO<sup>1</sup>, Teapa; BRITISH HONDURAS, R. Hondo (*Blancaneaux*).—AMAZONS, Santarem; BRAZIL, St. Paulo<sup>1</sup>.

"♂. Subovate, slightly convex, rather closely and visibly punctate, black with pale yellow markings; head yellow; thorax with the sides (very broadly) and front yellow, the lateral patch angulated on its inner edge; elytra each with two moderately large pale yellow spots—one discoidal, subtransverse, at the middle, one subapical, free, nearer the external margin. L. 1-1½ lin. (2-3 millim.).—♀. Head and front of thorax black."

A single example from British Honduras in our possession agrees entirely with Crotch's type, which is from Reiche's collection. There is a second example in Crotch's collection, from Santarem on the Upper Amazon.

vii. *Elytra with one yellow spot.*

19. **Hyperaspis centralis.** (Tab. XI. fig. 6.)

*Hyperaspis centralis*, Muls. Spec. Col. Trim. sécur. p. 685<sup>1</sup>; Crotch, Rev. Coccin. p. 229<sup>2</sup>.

*Hab.* MEXICO<sup>1,2</sup>, Toxpam (*Sallé*), Cuernavaca in Morelos, Jalapa, Cordova, Oaxaca, Tapachula in Chiapas (*Höge*); BRITISH HONDURAS, Belize (*Blancaneaux*); GUATEMALA, Guatemala city (*Salvin, Champion*), El Reposo, Zapote, Calderas, Chiacam, Chacoj (*Champion*); PANAMA, Bugaba, David, Caldera, Taboga I. (*Champion*).