

SELANDRIA.

Selandria, Leach, Zool. Misc. iii. p. 126 (1817); Hart. Blattw. p. 263 (1837).

Twenty-five species are referred by Kirby (List of Hymenoptera, p. 187) to this genus—fourteen from Europe, two from North America, three from Brazil, and six from Mexico; but two of these latter I have transferred to *Strongylogaster* *.

1. *Selandria luteola*.

Nigra, abdomine pedibusque rufo-luteis, ore albido, coxis, trochanteribus, femoribus basi, tibiis posticarum apice tarsisque nigris; alis fumatis, costa et stigmate fuscis.

Long. 4–5 millim.

Hab. PANAMA, Volcan de Chiriqui 3000 to 4000 feet (*Champion*).

The antennæ are stout, as long as the abdomen, the third joint is much longer than the fourth; the eyes converge on the lower side in front, and reach to the base of the mandibles; the frontal area is obsolete, the antennæ are widely separated, the lateral sutures on the vertex wide, but not deep; the antennal foveæ round, large; clypeus hairy, transverse, palpi white; the head retreats behind the eyes. The first transverse cubital nervure is subobsolete; the second cubital cellule is a very little longer than the third, which is longer than the fourth; the first recurrent nervure is received a little in front of the middle of cellule; the transverse median nervure is received a little in front of the apical third of cellule; the transverse radial nervure is largely bullated in the middle. The apex of coxæ is testaceous; the black on the femora reaches almost to the middle; behind the tibiæ are more or less fuscous; the anal appendages are black, pilose.

2. *Selandria curialis*.

Selandria curialis, Cresson, Trans. Amer. Ent. Soc. viii. p. 15¹.

Hab. MEXICO¹.

3. *Selandria flavipes*.

Selandria flavipes, Norton, Proc. Bost. Soc. N. H. viii. 1861, p. 222. 12; Trans. Amer. Ent. Soc. i. p. 258; iv. p. 85; Catal. p. 121¹.

Selandria flavipes, Cresson, Trans. Amer. Ent. Soc. viii. p. 60².

Hab. NORTH AMERICA.—MEXICO¹.

4. *Selandria glabra*.

Selandria glabra, Kirby, List of Hymen. i. p. 190, t. 8. fig. 23¹.

Hab. MEXICO, Orizaba¹.

* *S. diversipes*, Kirby, and *S. inconspicua*, Kirby.