

3. *Loboceras saussurii*.

Testaceum, pilosum, antennis (basi excepta), vertice, femoribus, tarsis abdominisque apice nigris; coxis, trochanteribus tibiisque albidis; alis flavo-hyalinis, apice fumatis, costa nigra, stigmatibus testaceo.

Long. 8 millim.

Hab. MEXICO, Orizaba (*de Saussure*).

The sutures on vertex are broad and deep, and there is a narrower one behind the ocelli; the former run down the sides of the frontal area and unite round its lower border, enclosing it completely, but at the angles of the area they are not so deep. The clypeus is very slightly incised at the apex; the mandibles are piceous at the apex. The first cubital cellule is small; the second is nearly double the length of the third, which is only a little longer than the first; the fourth is longer than the first three together. The curved first recurrent nervure is received close to the middle of the cellule; the second is interstitial, but it is more in the third cellule than in the second; the transverse median nervure is received a little in front of the middle of the cellule. The costa is black to near the commencement of the first cubital cellule; the apical cloud in the wings commences at the end of the stigma, and it curves back to the second transverse cubital nervure. The anterior femora are broadly white at the base. On the vertex the entire space bounded by the vertical and frontal sutures is black.

This species appears to be very closely related to *L. mexicanum*; but that has no black on the vertex. The costa is testaceous, like the stigma, and the tibiae are testaceous.

4. *Loboceras calcar*. (Tab. I. fig. 12.)

Cephalocera calcar, Norton, Trans. Amer. Ent. Soc. i. p. 51; Catal. p. 20 (♀)¹.

Hab. MEXICO¹.

This species differs from the others in having the first cubital cellule longer than the third, and in the transverse basal nervure being received quite close to the transverse basal.

5. *Loboceras mexicanum*.

Loboceras mexicanum, Kirby, List of Hymen. i. p. 80, t. 6. f. 4¹.

Hab. MEXICO, Orizaba¹.

6. *Loboceras xanthostigma*.

Testaceum, pilosum, antennis, capite (ore excepto), abdominis apice, tarsis tibiisque posterioribus nigris; alis flavo-hyalinis, apice fusco, costa et stigmatibus flavis.

Long. 7-8 millim.

Hab. PANAMA, Bugaba (*Champion*).

The face below and a little above the antennae is pallid yellow; the scape is more or