

Antennæ with the scape a little shorter than the third joint, the latter nearly equal in length to the fourth joint; the middle and apical joints slightly dilated at the apex. Head transverse in front, rugosely punctured, the face covered with long silvery-white hair; a keel down the centre of the front. Mesonotum transverse, shining; the middle lobe bearing large deep punctures, the lateral lobes slightly shagreened. Parapsidal furrows deep, wide; the furrow along the sides crenulated. Scutellum rugosely punctured. Pleuræ rugosely punctured, the punctures running into reticulations; the mesopleuræ in front in the middle impunctate. Metathorax coarsely rugosely punctured. Petiole shining impunctate, double the length of the metanotum. Pleuræ and metathorax densely covered with rather long white pile; a fuscous pile on the mesonotum. The petiole sparsely pilose; the apex of the abdomen above sparsely covered with short greyish pile. Coxæ thickly, the femora more sparsely, covered with a silvery-white pile; tibiæ and tarsi with a short thick fuscous pile. Hind coxæ somewhat coarsely punctured; tibiæ and tarsi covered with a few stiff bristles. Hind spurs reaching nearly to the middle of the metatarsus, the latter as long as the following three joints united. Claws apparently simple. The anterior legs are more or less obscure white in front. The transverse medial nervure is interstitial.

8. *Evania ornaticornis*. (Tab. XVII. fig. 20.)

Nigra, antennarum articulis 2^o-4^m apiceque petioli albis; fronte et facie punctatis; coxis posticis fere lævis; petiolo lævi; alis fere hyalinis. ♂ ♀.

Long. 7-8 millim.

Hab. PANAMA, Volcan de Chiriqui 4000 to 6000 feet (*Champion*).

Antennæ thick, slightly attenuated towards the apex; the scape as long as the third and fourth joints united; the third joint three times the length of the second, and a little longer than the fourth joint. Head rugosely punctured, the front a little depressed, a keel running down its centre from the ocelli, the space on either side of the keel above the antennæ being finely punctured or shagreened. Face almost transverse, keeled on either side. Mesonotum transverse in front; rugosely punctured, except the inner side of the lateral lobes. Parapsidal furrows moderately deep; there is no lateral furrow. Scutellum rugosely punctured. Pleuræ rugosely reticulated, except beneath the wings on the mesopleuræ, where there is a large shagreened space. Metathorax coarsely rugosely reticulated. Petiole impunctate, nearly twice the length of the metanotum. Hind coxæ indistinctly punctured, shining; hind spurs reaching nearly to the middle of the metatarsus, the latter as long as the following three joints united.

The male has the scape longer and thinner than the female; the third joint is longer compared to the fourth; and, if anything, the thorax is more closely punctured.