

distinct. Scutellum rugosely punctured. The lower part of the mesopleuræ and sternum strongly punctured, the upper part finely punctured, the oblique depression at the end obliquely striated. Metathorax rugosely reticulated. Petiole shining, impunctate, about one and a half times longer than the metanotum. Hind coxæ finely punctured; spurs reaching nearly to the middle of the metatarsus. The legs are rather long; the coxæ are obscure white, black at the base and apex; the trochanters are white, black at the apex, the posterior pair broadly so; the hind femora, tibiæ, and tarsi are black, the femora obscure brown in the middle. The mesonotum laterally and the metanotum in the middle are obscured with black.

2. *Radial and cubital nervures entirely obsolete.* (Species 12.)

12. **Evania nitida.** (Tab. XVIII. fig. 1, ♂.)

Rufa, lævis, antennis, metathorace postice, abdomine pedibusque posticis nigris, apice petioli facieque albis; alis hyalinis, apice fere fumatis. ♂.

Long. 4-5 millim.

*Hab.* PANAMA, Bugaba (*Champion*).

Antennæ issuing from near the middle of the eyes; scape scarcely so long as the second and third joints united; the third joint about three times the length of the second and slightly shorter than the fourth joint. Face projecting in the middle, obscurely punctured; the vertex more closely and strongly punctured. Pro- and mesothorax shining, impunctate; the middle lobe raised; the parapsidal furrows wide, deep; mesopleuræ deeply obliquely excavated, very shining. Scutellum shining, impunctate, the space in front of it crenulated. Metathorax rugosely punctured; the apex with an oblique rounded slope, black; the metapleuræ hollowed, shining, obliquely striated round the edges, the centre being smooth. The thorax is almost glabrous. Petiole more than double the length of the metanotum, shining, impunctate. The four anterior legs are pale reddish-testaceous, the middle femora blackish above. The hind legs are black, except the greater part of the coxæ and the base of the trochanters; they are sparsely covered with white hairs, the tarsi densely so. The long spur on the tibiæ is more than one third of the length of the metatarsus, the latter being a little shorter than the other joints united. Tarsi longer than the tibiæ, the joints elongated, the basal pair as long as the tibiæ.

b. *Parapsidal furrows obsolete. Cubital nervure obsolete beyond the first cubital cellule.* (Species 13.)

13. **Evania trochanterata.** (Tab. XVII. fig. 23, ♀.)

Nigra, rugosa, punctata; pro- et mesothorace rubris, antennarum articulis 2<sup>o</sup>-4<sup>m</sup>, trochanteribus, spinis petiolique dimidio apicali albis; petiolo lævi; alis hyalinis. ♀.

Long. 6.5 millim.

*Hab.* PANAMA, Volcan de Chiriqui 4000 to 6000 feet (*Champion*).