

Antennæ with the scape nearly as long as the third and fourth joints united; the third joint longer than the fourth, and nearly three times the length of the second joint. Face densely covered with stiff silvery-white pubescence, rugosely punctured, nearly transverse. Front densely pilose, the pile not so long as that on the face but closer, keeled down the centre, rugosely punctured; the vertex is not so strongly punctured, and has the punctures more widely separated. Collar almost transverse, but projecting a little in the middle. Mesonotum shining, the middle lobe marked with large distinctly separated punctures; the lateral lobes impunctate, a wide shallow furrow down the centre; parapsidal furrows narrow, moderately deep. Scutellum punctured. Pleuræ rugosely punctured, the punctures running into reticulations; a large shining impunctate space below the fore wings, and a smaller one below the hind wings. Metathorax rugosely punctured. Petiole double the length of the metanotum, smooth, shining, impunctate. Hind coxæ punctured; spurs reaching to the middle of the metatarsus, the latter scarcely so long as the following three joints united. The four anterior legs are more or less obscure brownish in front; the anterior trochanters are almost entirely white, the posterior pair having only the basal half white.

B. *Radial and cubital nervures obsolete. Antennæ with the third joint not more than double the length of the fourth; the other joints not much shorter than the third. Parapsidal furrows obsolete. Metatarsus generally longer than all the other joints united. Body more or less red.* (Species 14-18.)

#### 14. **Evania ocellaria.**

*Evania ocellaria*, Schletterer, Verh. zool.-bot. Ges. Wien, xxxvi. p. 233 (1886)<sup>1</sup>.

*Hab.* MEXICO, Orizaba (*Bilimek*<sup>1</sup>).

#### 15. **Evania rugosa.** (Tab. XVII. fig. 25, ♀.)

Nigra, pro- et mesonotum rufis, rugosis; petiolo punctato; alis hyalinis. ♀.  
Long. fere 6 millim.

*Hab.* GUATEMALA, Dueñas (*Champion*).

Antennæ with the scape scarcely so long as the following three joints united; the second joint a little longer than the fourth, and shorter than the third. Head rugosely punctured throughout, the punctures coarsest on the face, sparsely covered with glistening white hair. Thorax above coarsely rugosely punctured; the pro- and mesopleuræ shining, punctured only on the lower side. Metapleuræ reticulated, the reticulations elongated in front. Petiole one half longer than the metanotum, shining, sparsely punctured, and sparsely pilose. Hind coxæ impunctate; spurs reaching to the middle of the metatarsus, the latter longer than all the other joints united.