

## DRYINUS.

*Dryinus*, Latreille, Hist. Nat. Ins. xiii. p. 228 (1804) [sec. Förster, Hymen. Stud. ii. p. 90 (1856)].

This genus is apparently more numerous represented in tropical than in temperate regions.

The group of *D. MACULICORNIS*.

*Vertex depressed; occiput concave; ocelli in a triangle; eyes converging in front; third joint of the antennæ double the length of the first; third joint of the anterior tarsi double the length of the fourth; prothorax above bluntly keeled, longer than the mesothorax; parapsidal furrows obsolete; body rather long, narrow.*

1. ***Dryinus maculicornis***. (Tab. XIX. fig. 5.)

Niger, antennis testaceis cum articulis 5<sup>o</sup>-7<sup>m</sup> nigris, clypeo mandibulisque albis, tarsis abdominisque apice testaceis. ♂ ♀.

Long. 6 millim.

*Hab.* PANAMA, Bugaba 800 to 1500 feet, Volcan de Chiriqui 2000 to 3000 feet (*Champion*).

Antennæ with the apical five joints of the flagellum thicker than the basal joints; joints 7-9 brownish beneath; the basal joint white on the lower side, fuscous above. Head closely punctured, covered (especially along the eyes) with long white hair; the clypeus in the middle and the tips of the mandibles piceous; palpi fuscous. Thorax covered with rather long white hair; the pro- and mesothorax closely and somewhat strongly punctured, the former laterally, on the lower side, finely, longitudinally aciculated; metanotum coarsely, longitudinally striolated. Abdomen scarcely so long as the thorax; the basal three segments finely aciculate, the apical segment almost impunctate. The legs are sparsely covered with white hair; more or less black above, the front part white at the apex, the femora and tibiæ more or less brownish beneath; the coxæ and anterior trochanters are obscure brownish-testaceous. The wings are milky-white at the extreme base between the transverse basal nervure and the base of the stigma (which is white), and fusco-hyaline in front of the stigma.

2. ***Dryinus alticola***. (Tab. XIX. fig. 6.)

Niger, antennarum articulo 5<sup>o</sup> tarsisque testaceis; alis fuscis, fascia medio apiceque hyalinis. ♀.

Long. fere 6 millim.

*Hab.* PANAMA, Volcan de Chiriqui 4000 to 6000 feet (*Champion*).

Head closely, longitudinally, and rather strongly punctured; vertex scarcely depressed; a narrow keel runs down from the ocelli; palpi whitish. Prothorax finely longitudinally striated; the posterior part raised into a rather long hump, which is rather sharp in the centre. Mesothorax closely and rather strongly punctured;