

The group of *D. CHIRIQUENSIS*.

Head as in D. melanocephalus; prothorax scarcely so long as the mesothorax, quadrate (but, if anything, longer than broad), flat above, the sides straight; parapsidal furrows obsolete; metathorax with a rounded gradual slope.

Chiefly distinguished from the *D. melanocephalus* section by the obsolete parapsidal furrows, the longer and straighter prothorax, and the radial nervure being almost obsolete from a little beyond the apex of the stigma. The antennæ are missing from the example of the single species I include in this group.

6. *Dryinus chiriquensis*. (Tab. XIX. fig. 9.)

Niger, mandibulis, basi et apice abdominis pedibusque rufo-testaceis, geniculis, coxis anterioribus tarsisque pallidis, apice femorum posticorum nigris. ♂.

Long. 5 millim.

Hab. PANAMA, Volcan de Chiriqui 2000 to 3000 feet (*Champion*).

Head wide, longitudinally striolate-reticulate; sparsely covered with white hair. Prothorax somewhat strongly punctured and covered with long white hair; mesothorax sparsely pilose, almost impunctate; metathorax punctured, with six small upper and three large lower areæ. Abdomen a little shorter than the thorax, shining, impunctate. Wings in greater part hyaline; there is a narrow smoky line on the transverse basal and humeral nervures, and a broad stripe at the stigma (which is white except at the apex), and the apex is also slightly clouded; the radial nervure does not extend much beyond the stigma.

The group of *D. ALBITARSIS*.

Vertex not depressed; occiput short, convex; eyes parallel; third joint of the antennæ shorter than the first and about one fourth longer than the second; prothorax shorter than the mesothorax.

7. *Dryinus albitarsis*. (Tab. XIX. fig. 10.)

Niger, ore, antennis, coxis, trochanteribus, femoribus tibiisque anticis rufo-testaceis, tarsis pallidis; alis hyalinis, fusco-bifasciatis, stigmatibus nigro, cum basi alba. ♂.

Long. fere 3 millim.

Hab. GUATEMALA, Panajachel 5000 feet (*Champion*).

Antennæ stout, longer than the head and thorax united; almost glabrous. Head finely rugosely punctured, sparsely clothed with short white hairs; eyes margined. Prothorax rugosely punctured (more coarsely so than the head), the pronotum behind not so strongly, and more shining than the rest; mesonotum aciculated; scutellum shining, impunctate; metathorax longitudinally rugosely punctured, with a curved