

2. *Epyris coxalis*.

Niger, capite punctato, scapo antennarum, coxis, trochanteribus femorumque basi albis; alis fuscis, apice violaceis. ♂.
Long. fere 9 millim.

Hab. PANAMA, Bugaba (*Champion*).

Antennæ thick, densely pilose, the first joint a little longer than the third. Head shining, covered with shallow widely separated punctures; antennal depressions deep and wide; mandibles almost impunctate. Thorax sparsely punctured, the punctures very shallow. Parapsidal furrows not reaching to the scutellum; the transverse furrow in front of the latter wide and deep. Scutellum almost without punctures. Metanotum irregularly punctured and blistered; a stout keel runs down the centre and a wide and deep furrow down either side; the apex with a moderate slope; the sides transversely striated. Abdomen shining, the sides and ventral surface inclining to piceous. Femora and tibiæ sparsely, the tarsi densely, pilose.

3. *Epyris erythropoda*. (Tab. XIX. fig. 14.)

Niger, capite rugoso, antennis, ore, mandibulis, pedibus abdominisque apice rufis; alis hyalinis, cum fascia medio fumata; tegulis testaceis; coxis nigris. ♂.
Long. 10 millim.

Hab. PANAMA, Bugaba, Volcan de Chiriqui 3000 to 4000 feet (*Champion*).

Antennæ shorter than the thorax; the scape bare, shining, curved, longer than the third joint; the flagellum densely pilose, tapering perceptibly towards the apex; the third joint somewhat longer than the fourth. Head not much broader than the mesothorax, rather long, sparsely covered with long fuscous hairs, rugosely punctured; a shining furrow in the centre between the eyes; antennal tubercles large, reddish, a sharp projecting keel between them; mandibles with some large separated punctures, shining, the teeth black; palpi rather long, pilose, testaceous. Pro- and mesothorax bearing large punctures; the sternum and pleuræ more strongly punctured than the upper sides, except a shining impunctate space on the hinder part of the mesopleura; scutellum shining, finely but not closely punctured. The base of the metanotum bears some stout longitudinal keels; the rest of the flat posterior part is strongly transversely striolated, the two hinder parts being separated by a stout curved keel; the centre is depressed; the semiperpendicular apex is closely and strongly transversely punctured, except at the extreme top, the latter being excavated and the centre hollowed; the deep oblique depression at the base of the metapleuræ is shining and striated. Abdomen shining; the apex blunt, depressed in the centre, and very pilose, the hairs being fulvous. Legs densely pilose. Wings: radial nervure curved at the base; the recurrent nervure very faint, almost obsolete.