

Pelopæus assimilis, Dahlbom, Hymen. Eur. i. p. 23.

Pelopæus annulatus, Cresson, Proc. Ent. Soc. Phil. 1865, p. 135².

Hab. NORTH AMERICA.—MEXICO, Tampico¹, Cordova¹, Orizaba¹.—SOUTH AMERICA, Brazil, Chili, Patagonia; ANTILLES, Cuba².

✓ 3. **Pelopæus histrio.** (Tab. II. figg. 11; 11 a, ♂.)

Pelopæus histrio, Lepel. de St.-Fargeau, Hist. Nat. Ins., Hymén. iii. p. 316.

Pelopeus histrio, Saussure, Reise d. Novara, Hymen. p. 32¹.

Pelopæus bimaculatus, Lepel. de St.-Fargeau, loc. cit. p. 319.

Pelopæus fistularius, Dahlbom, Hymen. Eur. i. p. 23.

Hab. MEXICO¹, Atoyac in Vera Cruz (*Schumann*); GUATEMALA, El Reposo, Pantaleon, San Gerónimo, Panzos in Vera Paz (*Champion*).—SOUTH AMERICA¹, Cayenne, Brazil.

CHALYBION.

Chalybion, Dahlbom, Hymen. Europ. i. p. 21 (1845).

i. *Wings dark-violaceous; body covered with black hair.*

✓ 1. **Chalybion cæruleum.**

Sphex cærulea, Linnæus, Syst. Nat. i. p. 941.

Pelopeus cæruleus, Saussure, Reise d. Novara, Hymen. p. 26¹.

Chalybion cæruleum, Patton, Proc. Bost. Soc. Nat. Hist. xx. p. 378.

Hab. NORTH AMERICA, Middle and Southern States¹.—MEXICO, Cordova¹, Jaral in Guanajuato (*Schumann*).

✓ 2. **Chalybion zimmermanni.** (Tab. II. figg. 12; 12 a, ♂.)

Pelopæus zimmermanni, Dahlbom, Hymen. Eur. i. p. 22¹; Patton, Proc. Bost. Soc. Nat. Hist. xx. p. 379.

Pelopeus zimmermanni, Saussure, Reise d. Novara, Hymen. p. 26².

Hab. NORTH AMERICA, Carolina¹.—MEXICO, Ventanas (*Forrer*), Cordova², Michoacan², Valladolid in Yucatan (*Gaumer*).

C. (Pelopeus) californicum, Saussure (Reise d. Novara, Hymen. p. 26), is no doubt identical with this species.

ii. *Wings pellucid, smoky at the apex; body covered with cinereous hair.*

✓ 3. **Chalybion aztecum.**

Pelopeus aztecus, Saussure, Reise d. Novara, Hymen. p. 26¹.

Hab. MEXICO, Atoyac in Vera Cruz (*Schumann*), Tampico¹, Jalapa¹, Michoacan¹, Teapa in Tabasco (*H. H. Smith*).

BIOL. CENTR.-AMER., Hymenopt., Vol. II., November, 1888.

EE