

Hoplissus fuscus, Taschenberg, Zeitschr. f. Naturw. xlv. p. 368².

Gorytes fuscus, Handl. Sitz. der k. Akad. der Wissensch. Wien, xcvi. p. 377³.

Hab. NORTH AMERICA, Connecticut³.—MEXICO, Cordova³, Orizaba³, Temax in Northern Yucatan (*Gaumer*); GUATEMALA, San Joaquin in Vera Paz (*Champion*).—AMAZONS, Ega¹; BRAZIL, Rio Janeiro².

4. **Gorytes robustus.**

Gorytes robustus, Handl. Sitz. der k. Akad. der Wissensch. Wien, xcvi. p. 380¹.

Hab. MEXICO, Tampico¹.—SOUTH AMERICA, Brazil¹.

5. **Gorytes cayennensis.**

Hoplissus cayennensis, Spinola, Ann. Soc. Ent. Fr. x. p. 116¹.

Gorytes cayennensis, Handl. Sitz. der k. Akad. der Wissensch. Wien, xcvi. p. 385².

Gorytes sericatus, Smith, Cat. Hym. Ins. iv. p. 363³.

Hoplissus anthracipennellus, Taschenberg, Zeitschr. f. Naturw. xlv. p. 366⁴.

Hab. MEXICO, Orizaba (*Bilimek*²), Cordova (*Saussure*²).—SOUTH AMERICA, Colombia^{2 4}, Cayenne¹, Brazil^{3 4}.

6. **Gorytes maculipennis.** (Tab. V. figg. 12; 12 *a*, head from in front; 12 *b*, antenna.)

Niger, nitidus; thorace superne sparse punctato; scapo antennarum, margine clypei, orbitis oculorum, margine pronoti, scutello, fascia metanoti, maculis 2 in segmento mediali, tegulis fasciisque 4 abdominis, flavis; alis fere hyalinis, fasciis 3 fuscis.

Long. 7 millim.

Hab. PANAMA, Bugaba (*Champion*).

Antennæ stout, subclavate; the scape elongate, narrow, yellow beneath; the basal three-fourths of the flagellum brownish beneath; the third joint distinctly longer than the fourth. Head not much broader than the thorax, shining, impunctate, covered with a close fuscous pile; the clypeus with silvery hair; eyes almost parallel; hinder ocelli separated by a little more than the distance which separates them from the eyes; front not depressed, and without keel or furrow beyond a very thin indistinct furrow over the antennæ; clypeus short, broadly rounded, depressed at the base, fringed all round with silvery hair; mandibles piceous, blackish at the tips. Thorax above sparsely covered with a fuscous pile, the metapleuræ and sternum with a thicker and longer dull silvery pile, shining; the mesonotum and apex of the median segment covered with shallow punctures; the median area in the middle extends to near the centre of the median segment; the twelve keels gradually decrease from the centre to the sides, and are not distinctly bordered; the middle furrow is narrow, but distinct and continuous; mesosternal furrow narrow, but distinct. Abdomen a little longer and narrower than the thorax; the petiole and first segment impunctate; the other