

covered with stiff black hairs; the rest of the ventral surface is thickly covered with depressed pale ochreous hairs, which completely hide the sculpture. The coxæ and femora are covered with long, soft, pale hairs; the tarsal hairs are stiffer and black.

The following are the females of this section unknown to me:—

91. **Sphærophthalma canina.**

Mutilla canina, Smith, Cat. Hymen. iii. p. 58¹.

Sphærophthalma canina, Blake, Trans. Am. Ent. Soc. xiii. p. 215².

Hab. MEXICO^{1 2}.

92. **Sphærophthalma combusta.**

Mutilla combusta, Smith, Cat. Hymen. iii. p. 59¹.

Sphærophthalma combusta, Blake, Trans. Am. Ent. Soc. xiii. p. 216².

Hab. MEXICO^{1 2}.

93. **Sphærophthalma vestita.**

Mutilla vestita, Lepel. de St.-Farg. Nat. Hist. Ins., Hymén. iii. p. 634¹.

Sphærophthalma vestita, Blake, Trans. Am. Ent. Soc. xiii. p. 218².

Hab. MEXICO^{1 2}.

94. **Sphærophthalma verticalis.**

Mutilla verticalis, Smith, Cat. Hymen. Ins. iii. p. 63¹.

Sphærophthalma verticalis, Blake, Trans. Am. Ent. Soc. xiii. p. 221².

Hab. MEXICO^{1 2}.

95. **Sphærophthalma pulchra.**

Mutilla pulchra, Smith, Cat. Hymen. Ins. iii. p. 62¹.

Sphærophthalma pulchra, Blake, Trans. Am. Ent. Soc. xiii. p. 222².

Hab. MEXICO^{1 2}.

96. **Sphærophthalma antiguensis.**

Mutilla antiguensis, Fabr. Ent. Syst. ii. p. 367¹; Syst. Piez. p. 429².

Sphærophthalma antiguensis, Blake, Trans. Am. Ent. Soc. xiii. p. 221 (♀, ♂)³.

Hab. MEXICO³.—ANTILLES, Antigua^{1 2 3}.

97. **Sphærophthalma klugi.**

Mutilla klugii, Gray, in Griffith's Anim. Kingd., Ins. ii. p. 516, t. 71. fig. 2¹; Lepel. de St.-Farg. Nat. Hist. Ins., Hymén. iii. p. 596².

Hab. ? MEXICO.—SOUTH AMERICA¹.

I introduce this species on the authority of a MS. note of the late Mr. F. Smith, in which he gives "Mexico" as a habitat.