

externo quoque nigricantibus: subtus fuscis, lineis tribus communibus albis, una discali introrsum fusco limbata et marginem posticarum internum bene angulata, duabus submarginalibus interiore extrorsum fusco (ad marginem posticarum internum nigro) limbata, lobulo anali nigro, area juxta eum introrsum rubra, macula magna rubra inter ramos medianos puncto nigro, margine externo anguste nigro introrsum limbato.

♀ mari similis, colore cæruleo alarum paullo dilutiore.

*Hab.* NICARAGUA, Chontales (*Belt*); COSTA RICA, Cache, Rio Sucio (*Rogers*); PANAMA, Volcan de Chiriqui (*Ribbe*).—VENEZUELA; ECUADOR<sup>1 2</sup>.

A well-marked species, having no very near allies, of which we have a considerable series of specimens ranging from Nicaragua to the State of Panama.

#### 14. T. ODINUS section.

##### 139. *Thecla odinus*, sp. n. (Tab. LVI. figg. 20, 21 ♀.)

Alis fusco-nigris; anticis cellulæ basi et infra eam ad marginem internum et posticis interne cæruleis, his linea submarginali ad angulum analem quoque cærulea: subtus rufescenti-brunneis; anticis ad marginem externum ferrugineis, linea discali communi argenteo-cærulea in posticis perfracta, lineola altera ad cellulæ alarum finem coloris ejusdem, lineis duabus submarginalibus pallidis interiore argenteo-cæruleo atomata, lobulo anali et macula inter ramos medianos nigris, inter eos macula argentea omnibus ferrugineo introrsum limbatis, margine interno nigro.

♂ adhuc ignotus.

*Hab.* PANAMA, Volcan de Chiriqui (*Champion*).

We have but a single female specimen of this pretty species, taken by Mr. Champion on the Volcan de Chiriqui, at an altitude of between 2500 to 4000 feet above the sea.

#### 15. T. MÆONIS section.

##### 140. *Thecla mæonis*, sp. n. (Tab. LVI. figg. 22, 23 ♂.)

Alis fusco-nigris; anticis cellula et infra eam ad angulum analem et posticis præter costam et marginem externum læte cæruleis, ciliis rufescentibus: subtus ferrugineis, lineis duabus communibus transversis albis, una discali rufescente introrsum limbata, altera submarginali extrorsum rufo (ad marginem internum nigro) marginata, lobulo anali et puncto inter ramos medianos nigris introrsum rufo late limbatis, macula inter eos quoque nigra albo atomata.

♀ nobis ignota.

*Hab.* MEXICO, Jalapa (*W. Schaus*); GUATEMALA, San Gerónimo (*F. D. G. & O. S., Champion*).

We have two specimens (one of which we figure) of this species, both from San Gerónimo in Guatemala. These agree with Mexican specimens in Mr. Schaus's collection. The species has no near ally with which we are acquainted.

#### 16. T. PARON section.

##### 141. *Thecla paron*, sp. n. (Tab. LVI. figg. 24, 25 ♂.)

Alis fusco-nigris; anticis infra cellulam ad marginem internum cæruleo atomatis, macula indistincta nigra ad