

**Euptychia antinoe** (I. p. 86).*Euptychia antinoe*, Staud. Exot. Schmett. p. 225, t. 80<sup>3</sup>.

To the localities given, add:—MEXICO, Teapa (*H. H. Smith*); HONDURAS<sup>3</sup>, San Pedro Sula (*Wittkugel*); PANAMA, Bugaba, Tolé (*Champion*).

**Euptychia camerta** (I. p. 86).*Euptychia camerta*, Staud. Exot. Schmett. p. 226, t. 81 (*E. hermes*)<sup>12</sup>.

Of this common species we have received specimens from a large number of additional localities; but as these latter add nothing to the general distribution of the insect, it is unnecessary to enumerate them here.

**Euptychia ebusa** (I. p. 88).

To the localities given, add:—PANAMA, Bugaba, Volcan de Chiriqui, David (*Champion*).

Sent in numbers by Mr. Champion from Chiriqui.

**Euptychia philodice** (I. p. 90).

To the localities given, add:—PANAMA, Volcan de Chiriqui 8000 feet (*Champion*).

**Euptychia nelsoni** (I. p. 91). (Tab. CVII. figg. 7, 8, ♂.)

We now give a figure of this very distinct species.

**Euptychia argentella** (I. p. 91).*Euptychia argentella*, Staud. Exot. Schmett. p. 227, t. 81<sup>3</sup>.

To the localities given, add:—MEXICO, Omilteme, Xucumanatlan, and Xautipa, in Guerrero, Orizaba (*Elwes*); GUATEMALA<sup>3</sup>, Volcan de Atitlan, San Lucas Toliman (*Champion*).

We have six specimens of a variety of this species which are much redder than the typical form; these are from San Lucas and Cerro Zunil in Guatemala, whence we also have normally-coloured individuals.

31 (A). **Euptychia clinas**. (Tab. CVII. figg. 9, 10, ♂.)*Euptychia clinas*, Godm. & Salv. Ann. & Mag. Nat. Hist. (6) iii. p. 352 (1889)<sup>1</sup>.

Alis anticis ad apicem obtusis, margine externo concavo; posticis margine externo dentato, angulo anali bene producto, fuscis; anticis ad costam et ad marginem externum saturatioribus, posticis ad angulum apicalem quoque obscurioribus maculis tribus submarginalibus nigricantibus notatis: subtus pallidioribus et paullo rufescentioribus, lineis tribus undulatis communibus transfasciatis, una per cellulas, secunda discali, tertia submarginali; posticis maculis duabus ad angulum analem rotundis, altera sagittiformi inter ramos medianos, argenteis, ocellis duobus ad medium marginis externi nigris fulvo circumcinctis, maculis argenteis bipupillatis.

♀ nobis ignota.