

NOTE.—Of the three unidentified species of *Neonympha* placed by us at the end of the genus *Euptychia* (anteà, i. p. 94), two have now been identified:—*N. epinephile*, Feld. = *Paramecera xicaque*, Reak. (anteà, i. p. 101); *N. pompilia*, Feld. = *Euptychia pieria*, Butl. (anteà, i. p. 79).

TAYGETIS (I. p. 94).

Taygetis armillata (I. p. 96).

To the localities given, add:—PANAMA, Veraguas (*Arcé*).

Taygetis virgilia (I. p. 97).

Taygetis virgilia, var. *rufomarginata*, Staud. Exot. Schmett. p. 235, t. 85 (1888)⁴.

To the localities given, add:—MEXICO, Jalapa (*coll. Schaus*), Atoyac (*H. H. Smith*); GUATEMALA, Volcan de Santa Maria (*Richardson*), Cerro Zunil (*Champion*); HONDURAS, San Pedro Sula (*Wittkugel*); NICARAGUA, Matagalpa (*Richardson*); PANAMA, Bugaba, Volcan de Chiriqui (*Champion*), Veraguas (*Arcé*).

Of this species we have now seen many more specimens from within our limits. The var. *rufomarginata* is the commoner form from Guatemala to Panama.

Taygetis nympha (I. p. 97).

To the localities given, add:—GUATEMALA, San Isidro (*Champion*).

Two specimens of this species have been sent us by Mr. Champion from the Pacific slope of Guatemala.

Taygetis andromeda (I. p. 98).

Taygetis salvini, Staud. Exot. Schmett. p. 236, t. 85 (1888)¹⁰.

To the localities given, add:—MEXICO, Jalapa (*coll. Schaus*), Cuesta de Misantla (*M. Trujillo*), Atoyac, Teapa (*H. H. Smith*); BRITISH HONDURAS, Rio Hondo (*Blancaneaux*); GUATEMALA, Volcan de Santa Maria (*Richardson*); NICARAGUA, Matagalpa (*Richardson*); COSTA RICA, Santa Clara Valley (*Zurcher*); PANAMA, Bugaba (*Champion*).

With our very extensive series of this extremely variable species we are quite unable to separate *T. salvini* as distinct. Dr. Staudinger's type of the latter was from Chiriqui¹⁰.

Taygetis keneza (I. p. 99).

To the localities given, add:—MEXICO, Rio Papagaio and Dos Arroyos in Guerrero (*H. H. Smith*); PANAMA, Bugaba (*Champion*).

Sent in abundance by Mr. Champion from Bugaba.