

indistinctis alis transeuntibus, una per cellulas, secunda ultra eas, in posticis latiore, recta et rubescentiore, tertia submarginali, in anticis extrorsum ferrugineo limbata, posticis punctis quatuor submarginalibus nigris albo pupillatis.

*Hab.* COSTA RICA (*Underwood* <sup>1</sup>).

We are unacquainted with this insect, and have taken our description from Mr. Grose-Smith's work. It appears to be most nearly allied to *P. hulda*, Butl.

**Pedaliodes triaria** (I. p. 105).

To the localities given, add:—PANAMA, Volcan de Chiriqui 8000 feet (*Champion*).

GYROCHEILUS (I. p. 106).

**Gyrocheilus patrobas** (I. p. 106).

*Gyrocheilus patrobas*, Staud. Exot. Schmett. p. 234, t. 84 (♀) <sup>4</sup>.

*Geirocheilus tritonia*, W. H. Edwards, Trans. Am. Ent. Soc. v. p. 18 (1874) <sup>5</sup>; Butt. N. Am., *Geirocheilus*, i. ff. 1-4 (♂ ♀) <sup>6</sup>.

To the localities given, add:—NORTH AMERICA, Arizona <sup>5 6</sup>. — MEXICO <sup>4</sup>, Northern Sonora (*Morrison*), Durango city (*Becker*), Jalapa (*coll. Schaus*).

Morrison and Becker have sent us numerous examples from Northern Mexico agreeing with Mr. Edwards's figures of *G. tritonia*, and these insects are inseparable from *G. patrobas* (Hew.).

OXEOSCHISTUS (I. p. 106).

**Oxeoschistus hilarus** (I. p. 107).

To the localities given, add:—MEXICO, Xautipa and Omilteme in Guerrero (*H. H. Smith*).

**Oxeoschistus euryphile** (I. p. 107).

To the localities given, add:—PANAMA, Volcan de Chiriqui 8000 feet (*Champion*).

**Oxeoschistus tauropolis** (I. p. 108).

To the localities given, add:—MEXICO, Cuesta de Misantla (*M. Trujillo*), Jalapa (*coll. Schaus*), Orizaba (*Elwes*); GUATEMALA, Chiacam and Sabo in Vera Paz, Volcan de Atitlan (*Champion*).

**Oxeoschistus puerta** (I. p. 109).

*Oxeoschistus simplex*, Staud. Exot. Schmett. p. 234, t. 84 (♂) <sup>4</sup>.

Dr. Staudinger has figured a specimen of this species from Colombia under the name of *O. simplex*.