

Caligo memnon (I. p. 133).

To the localities given, add:—MEXICO, Teapa (*H. H. Smith*); GUATEMALA, Volcan de Santa Maria (*Richardson*), La Tinta and Panima in Vera Paz, Volcan de Atitlan (*Champion*); HONDURAS, San Pedro Sula (*Wittkugel*); PANAMA, Bugaba, Tolé (*Champion*).

Caligo atreus (I. p. 135).

Caligo atreus, Staud. Exot. Schmett. p. 215, t. 75 (♂) (1888)⁵.

To the localities given, add:—COSTA RICA, Santa Clara Valley (*Zurcher*); PANAMA, Bugaba (*Champion*).

Caligo uranus (I. p. 136).

To the localities given, add:—MEXICO, Cuesta de Misantla (*M. Trujillo*), Teapa (*H. H. Smith*); GUATEMALA, Cubilguitz and Telemán in Vera Paz (*Champion*).

ERYPHANIS (I. p. 136).

Eryphanis wardi (I. p. 137).

To the localities given, add:—PANAMA, Bugaba (*Champion*).

Eryphanis æsacus (I. p. 137).

Eryphanis æsacus, Staud. Exot. Schmett. p. 217, t. 76 (♂) (1888)³.

To the localities given, add:—GUATEMALA, Cahabon (*Champion*).

NAROPE (I. p. 139).

2. Narope anartes.

Narope anartes, Hew. Boliv. Butt. p. 9 (1874)¹.

Narope sarastro, Staud. Exot. Schmett. p. 218, t. 76².

Alis rufo-fulvis, anticis falcatis, apicem versus cum margine externo anguste, et posticis ad apicem, nigro-fuscis: subtus pallidioribus, fusco et brunneo irroratis, fascia communi ab apice anticarum ad medium marginis interni posticarum sordide ochracea, in posticis utrinque fusco-nigro marginatis.

Hab. COSTA RICA, Cartago (*Underwood, in coll. Druce*).—COLOMBIA, Cauca Valley²; BOLIVIA¹; PERU; ARGENTINE REPUBLIC.

Mr. Druce has a single specimen of this widely-distributed insect, sent him by Mr. Underwood in a collection of moths from Costa Rica. It agrees with others from the Cauca Valley and Bolivia, purchased from Dr. Staudinger under the name of *N. sarastro*, which we are unable to separate from *N. anartes*, Hewitson.

ACTINOTE (I. p. 140).

Actinote anteus (I. p. 141).

Acraea anteus, Staud. Exot. Schmett. p. 81, t. 32⁴.

To the localities given, add:—HONDURAS, San Pedro Sula (*Wittkugel*).