

**Synchloe hyperia** (I. p. 179).

*Chlosyne hippodrome*, Staud. Exot. Schmett. p. 95, t. 36<sup>7</sup>.

To the localities given, add :—MEXICO, Milpas in Durango (*Forrer*), Sierra Madre de Tepic, Tehuantepec (*Richardson*), Acaguizotla, Rio Papagaio (*H. H. Smith*), Coatepec (*Brooks*), San Lorenzo (*Trujillo*); PANAMA, San Lorenzo (*Champion*).

**Synchloe melanarge** (I. p. 179).

To the localities given, add :—MEXICO, Acaguizotla, Amula, La Venta, Rincon (*H. H. Smith*); NICARAGUA, Matagalpa (*Richardson*).

We have now received six specimens of this species from Mexico.

**Synchloe erodyle** (I. p. 180).

To the localities given, add :—GUATEMALA, Volcan de Santa Maria (*Richardson*); HONDURAS, San Pedro Sula (*Wittkugel*).

**Synchloe pæcile** (I. p. 180).

*Chlosyne pæcile*, Staud. Exot. Schmett. p. 95, t. 36<sup>3</sup>.

To the localities given, add :—PANAMA, Bugaba (*Champion*).

We omitted to note in our diagnosis of this species that the submarginal row of spots on the primaries are white, as in *S. erodyle*.

6 (A). **Synchloe eumeda**. (Tab. CVIII. figg. 1, 2, ♂.)

*Synchloë eumeda*, Godm. & Salv. Ann. & Mag. Nat. Hist. (6) xiv. p. 96<sup>1</sup>.

*S. pæcili* (formæ typicæ) similis, sed anticis maculis flavis in cellula et ea infra venam medianam evanescentibus, aliter fascia maculari ultra cellulam multo latiore, et punctis parvis submarginalibus flavis (nec albis); posticis fascia, venis divisa, ad cellulæ finem latissima, a costa ad marginem internum extendente aurantia, maculis quinque (nec quatuor) in serie submarginali rubris: subtus anticis maculis ad basin minoribus, aliis majoribus, omnibus flavis; posticis maculis nigris ad basin minoribus, maculis quinque submarginalibus rubris ut supra.

*Hab.* MEXICO, Mochitlan in Guerrero (*Baron*<sup>1</sup>).

We have four specimens of this species, which we have now ascertained are all of the male sex.

6 (B). **Synchloe dryope**. (Tab. CVIII. figg. 3, 4, ♂.)

*Synchloë dryope*, Godm. & Salv. Ann. & Mag. Nat. Hist. (6) xiv. p. 96<sup>1</sup>.

*S. eumedæ* similis, sed fascia maculari communi flava (nec aurantia): subtus posticis ad basin maculis majoribus et lineis distinctis formantibus.

*Hab.* MEXICO, Jalisco (*Richardson*<sup>1</sup>).

Of this species we have a single male specimen only; it is perhaps an extreme form of *S. eumeda*.